



Convergence Rates of Monotone Schemes for Conservation Laws for Data with Unbounded Total Variation

Ulrik Skre Fjordholm¹ · Kjetil Olsen Lye²

Received: 15 October 2020 / Revised: 13 January 2022 / Accepted: 7 February 2022 /
Published online: 21 March 2022
© The Author(s) 2022

Abstract

We prove convergence rates of monotone schemes for conservation laws for Hölder continuous initial data with unbounded total variation, provided that the Hölder exponent of the initial data is greater than $1/2$. For strictly Lip^+ stable monotone schemes, we prove convergence for any positive Hölder exponent. Numerical experiments are presented which verify the theory.

Keywords Hyperbolic conservation laws · Numerical methods · Convergence rate · Irregular data

1 Introduction

Consider the scalar hyperbolic conservation law

$$\begin{aligned}u_t + f(u)_x &= 0 \\ u(x, 0) &= u_0(x)\end{aligned}\tag{1}$$

where $f \in C^1(\mathbb{R})$ is the flux function and $u_0 \in L^1 \cap L^\infty(\mathbb{R})$ is the initial data. Equations of this form appear in a large number of applications, including scenarios where very irregular data is to be expected; we mention in particular flow in porous media [1, 2], turbulent flows (so-called “Burgulence”) [3–5], and advection with rough coefficients [6]. While the study of qualitative properties of “rough” solutions of (1) has been explored in detail (see e.g. [3–5]), the behavior of numerical methods for (1) has received much less attention. There exists convergence results for *linear* conservation laws with rough coefficients [6]. However, there are no known results for non-linear conservation laws with rough *initial data*.

✉ Ulrik Skre Fjordholm
ulriksf@math.uio.no

Kjetil Olsen Lye
kjetil.olsen.lye@sintef.no

¹ Department of Mathematics, University of Oslo, Postboks 1053 Blindern, 0316 Oslo, Norway

² Mathematics and Cybernetics, SINTEF, Pb. 124 Blindern, 0314 Oslo, Norway

The purpose of this paper is to study the convergence rate of monotone numerical methods for (1) in the presence of rough (say, piecewise Hölder continuous) initial data u_0 . As is to be expected, the convergence rate deteriorates with lower regularity. We demonstrate in several numerical experiments that our estimates are sharp, or close to being sharp.

1.1 Weak Solutions of Hyperbolic Conservation Laws

As is well-known, solutions of nonlinear hyperbolic Eq. (1) can develop shocks in finite time, making it necessary to interpret the equation in a weak manner. A weak solution of (1) is a function $u \in L^1 \cap L^\infty(\mathbb{R} \times \mathbb{R}_+)$ satisfying

$$\int_{\mathbb{R}_+} \int_{\mathbb{R}} u(x, t)\phi_x(x, t) + f(u(x, t))\phi(x, t) dx dt + \int_{\mathbb{R}} \phi(x, 0)u_0(x) dx = 0 \tag{2}$$

for all test functions $\phi \in C_c^\infty(\mathbb{R} \times \mathbb{R}_+)$. It is well-known that weak solutions are non-unique, so one introduces *entropy conditions* to single out the physically relevant solutions. Concretely, we say that $u \in L^1 \cap L^\infty(\mathbb{R} \times \mathbb{R}_+)$ is an entropy solution of (1), if for every pair of functions $\eta, q: \mathbb{R} \rightarrow \mathbb{R}$ where η is convex and $q' = \eta' f'$, it holds that

$$\eta(u)_t + q(u)_x \leq 0$$

in the sense of distributions. In particular, it is sufficient to impose the entropy condition with respect to the Kruzkov entropy pairs, given by

$$\eta(u, k) = |u - k|, \quad q(u, k) = \text{sign}(u - k)(f(u) - f(k)), \quad u \in \mathbb{R}$$

for all $k \in \mathbb{R}$. It was shown by Kruzkov (see e.g. [7] or [8, Proposition 2.10]) that entropy solutions of (1) are unique.

1.2 Finite Volume Methods for Conservation Laws

This section briefly describes the conventional approach of numerical approximation of conservation laws through finite volume and finite difference methods. For a complete review, one can consult e.g. [9].

We discretize the spatial domain \mathbb{R} by partitioning it into a collection of cells $C_i := [x_{i-1/2}, x_{i+1/2}) \subset \mathbb{R}$ with corresponding cell midpoints $x_i := \frac{x_{i+1/2} + x_{i-1/2}}{2}$. For simplicity we assume that our mesh is equidistant, that is,

$$x_{i+1/2} - x_{i-1/2} \equiv \Delta x \quad \forall i \in \mathbb{Z}$$

for some $\Delta x > 0$. We discretize time by equidistant points, that is, we choose $t^n = n\Delta t$ for $n \in \mathbb{N}_0$ for some $\Delta t > 0$.

For each cell C_i and each point in time t^n we let v_i^n be an approximation of the cell average of u at time t^n , $u_i^n \approx \int_{C_i} u(x, t^n) dx$ (here, $\int_C := \frac{1}{|C|} \int_C$, where $|C|$ is the Lebesgue measure of a Lebesgue set $C \subset \mathbb{R}$). This approximation is computed according to the finite volume scheme

$$\frac{v_i^{n+1} - v_i^n}{\Delta t} + \frac{F(v_i^n, v_{i+1}^n) - F(v_{i-1}^n, v_i^n)}{\Delta x} = 0 \tag{3}$$

$$v_i^0 = \int_{C_i} u_0(x) dx$$

where F is a *numerical flux function*. We furthermore assume the numerical flux function is consistent with f and locally Lipschitz continuous; more precisely, for every bounded set $K \subset \mathbb{R}$, there exists a constant $C_F > 0$ such that

$$|F(a, b) - f(a)| + |F(a, b) - f(b)| \leq C_F |b - a| \quad \forall a, b \in K. \tag{4}$$

We will frequently abuse notation and view grid functions $v \in \ell^1(\mathbb{Z})$ as an element of $L^1(\mathbb{R})$ under the inclusion $\ell^1(\mathbb{Z}) \hookrightarrow L^1(\mathbb{R})$ which maps $v \mapsto \sum_i v_i \mathbb{1}_{C_i}$.

2 A Modified Kuznetsov Lemma

Kuznetsov’s lemma [10] provides an explicit estimate of the difference between two (approximate) solutions of (1) in terms of their relative (Kruzkov) entropy. In this section we recall Kuznetsov’s lemma and prove a corollary which — as opposed to Kuznetsov’s original application of the lemma — does not depend on $\text{TV}(u_0)$ being bounded. We note that both Lemmas 1 and 2 can be generalized to multiple space dimensions (see e.g. [10] and [8, Section 4.3]). However, since we will only be able to handle one-dimensional equations in Sect. 3, we chose to stick to one dimension, for the sake of simplicity.

Fix now some final time $T > 0$. Kuznetsov’s lemma estimates approximation errors in the space

$$\mathcal{K} := \{u : \mathbb{R}_+ \rightarrow L^1(\mathbb{R}) \mid u \text{ and has right and left limits at all } (t, x) \in \mathbb{R}_+ \times \mathbb{R}\}.$$

For $u \in \mathcal{K}$ and $\sigma > 0$ we define the moduli of continuity

$$v_t(u, \sigma) = \sup \{ \|u(t + \tau) - u(t)\|_{L^1(\mathbb{R})} \mid 0 < \tau \leq \sigma \}, \quad v(u, \sigma) = \sup_{t \in [0, T]} v_t(u, \sigma).$$

Let $\omega \in C_c^\infty(\mathbb{R})$ be a standard mollifier, i.e. an even function satisfying $\text{supp } \omega \subset [-1, 1]$, $0 \leq \omega \leq 1$ and $\int_{\mathbb{R}} \omega dx = 1$. For $\epsilon > 0$ we define $\omega_\epsilon(x) = \frac{1}{\epsilon} \omega(\frac{x}{\epsilon})$. For $\epsilon, \epsilon_0 > 0$, define

$$\Omega(x, x', s, s') = \omega_{\epsilon_0}(s - s') \omega_\epsilon(x - x') \quad (x, x', s, s') \in \mathbb{R}^4.$$

For $\phi \in C_c^\infty(\mathbb{R} \times \mathbb{R}_+, \mathbb{R})$, $k \in \mathbb{R}$ and $u, v \in \mathcal{K}$ we set

$$\begin{aligned} \Lambda_T(u, \phi, k) &= \int_0^T \int_{\mathbb{R}} (|u - k| \phi_t + q(u, k) \phi_x) dx dt \\ &\quad - \int_{\mathbb{R}} |u(x, T) - k| \phi(x, T) dx + \int_{\mathbb{R}} |u(x, 0) - k| \phi(x, 0) dx, \\ \Lambda_{\epsilon, \epsilon_0}(u, v) &= \int_0^T \int_{\mathbb{R}} \Lambda_T(u, \Omega(\cdot, x', \cdot, s'), v(x', s')) dx' ds'. \end{aligned}$$

Lemma 1 (*Kuznetsov’s lemma [10]*) *Let $v \in \mathcal{K}$ and let w be an entropy solution of (1). If $0 < \epsilon_0 < T$ and $\epsilon > 0$, then*

$$\begin{aligned} \|v(\cdot, T) - w(\cdot, T)\|_{L^1(\mathbb{R})} &\leq \|v_0 - w_0\|_{L^1(\mathbb{R})} + \text{TV}(w_0) (2\epsilon + \epsilon_0 \|f\|_{\text{Lip}}) \\ &\quad + v(v, \epsilon_0) - \Lambda_{\epsilon, \epsilon_0}(v, w) \end{aligned}$$

where $v_0 = v(\cdot, 0)$ and $w_0 = w(\cdot, 0)$.

The following is a straightforward extension of [10, Lemma 4 and Theorem 4].

Lemma 2 Let $u_0 \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$ and let $v^{\Delta x}$ be the solution computed by a monotone finite volume scheme (3) with initial data $v_0^{\Delta x}$. Then

$$\|u(T) - v^{\Delta x}(T)\|_{L^1(\mathbb{R})} \leq 2\|u_0 - v_0^{\Delta x}\|_{L^1(\mathbb{R})} + \text{TV}(v_0^{\Delta x})(2\epsilon + \epsilon_0\|f\|_{\text{Lip}} + 2C_F \max(\epsilon_0, \Delta t)) + C \left(\frac{C_F \Delta x}{\epsilon} + \frac{\|f\|_{\text{Lip}} \Delta t}{\epsilon_0} \right) \sum_{n=0}^N \text{TV}(v^{\Delta x}(t^n)) \Delta t. \tag{5}$$

for any $T > 0, \epsilon > 0$ and $0 < \epsilon_0 < T$, for some $C > 0$ only depending on the choice of smoothing kernel ω .

Proof Let w be the entropy solution of (1) with $w_0 = v_0^{\Delta x}$. Then

$$\|u(T) - v^{\Delta x}(T)\|_{L^1(\mathbb{R})} \leq \|u(T) - w(T)\|_{L^1(\mathbb{R})} + \|w(T) - v^{\Delta x}(T)\|_{L^1(\mathbb{R})} \leq \|u_0 - v_0^{\Delta x}\|_{L^1(\mathbb{R})} + \|w(T) - v^{\Delta x}(T)\|_{L^1(\mathbb{R})}$$

by the stability of entropy solutions in $L^1(\mathbb{R})$ (see e.g. [11, Theorem 1] or [8, Proposition 2.10]). We estimate the second term using Lemma 1. For notational convenience, denote

$$\eta_i^n = |v_i^n - k| \quad \text{and} \quad q_i^n = q(v_i^n, k).$$

Without loss of generality we may assume that $T = t^{N+1}$ for some $N \in \mathbb{N}$. Then

$$\begin{aligned} -\Delta_T(v^{\Delta x}, \phi, k) &= -x \sum_{n=0}^N \sum_{i=-\infty}^{\infty} \int_{t^n}^{t^{n+1}} \int_{x_{i-1/2}}^{x_{i+1/2}} (\eta_i^n \phi_t + q_i^n \phi_x) dx dt \\ &\quad + \sum_{i=-\infty}^{\infty} \int_{x_{i-1/2}}^{x_{i+1/2}} \eta_i^{N+1} \phi(x, T) dx - \sum_{i=-\infty}^{\infty} \int_{x_{i-1/2}}^{x_{i+1/2}} \eta_i^0 \phi(x, 0) dx \\ &= -\sum_{n=0}^N \sum_{i=-\infty}^{\infty} \int_{x_{i-1/2}}^{x_{i+1/2}} \eta_i^n (\phi(x, t^{n+1}) - \phi(x, t^n)) dx \\ &\quad + \sum_{n=0}^N \sum_{i=-\infty}^{\infty} \int_{t^n}^{t^{n+1}} q_i^n (\phi(x_{i+1/2}, t) - \phi(x_{i-1/2}, t)) dt \\ &\quad + \sum_{i=-\infty}^{\infty} \int_{x_{i-1/2}}^{x_{i+1/2}} \eta_i^{N+1} \phi(x, T) dx - \sum_{i=-\infty}^{\infty} \int_{x_{i-1/2}}^{x_{i+1/2}} \eta_i^0 \phi(x, 0) dx \end{aligned}$$

(summation by parts)

$$\begin{aligned} &= \sum_{n=0}^N \sum_{i=-\infty}^{\infty} \int_{x_{i-1/2}}^{x_{i+1/2}} (\eta_i^{n+1} - \eta_i^n) \phi(x, t^{n+1}) dx \\ &\quad + \Delta x \sum_{n=0}^N \sum_{i=-\infty}^{\infty} \int_{t^n}^{t^{n+1}} (q_{i+1}^n - q_i^n) \phi(x_{i+1/2}, t) dt \end{aligned}$$

(set $\bar{\phi}_i^n := \frac{1}{\Delta x} \int_{x_{i-1/2}}^{x_{i+1/2}} \phi(x, t^{n+1}) dx$ and $\bar{\phi}_{i+1/2}^{n+1} := \frac{1}{\Delta t} \int_{t^n}^{t^{n+1}} \phi(x_{i+1/2}, t) dt$)

$$= \Delta x \sum_{n=0}^N \sum_{i=-\infty}^{\infty} (\eta_i^{n+1} - \eta_i^n) \bar{\phi}_i^n + \Delta t \sum_{n=0}^N \sum_{i=-\infty}^{\infty} (q_{i+1}^n - q_i^n) \bar{\phi}_{i+1/2}^{n+1}.$$

Let $Q_{i+1/2}^n := F(v_i^n \vee k, v_{i+1}^n \vee k) - F(v_i^n \wedge k, v_{i+1}^n \wedge k)$ be the Crandall–Majda numerical entropy flux, so that

$$\eta_i^{n+1} - \eta_i^n + \frac{\Delta t}{\Delta x} (Q_{i+1/2}^n - Q_{i-1/2}^n) \leq 0$$

(see e.g. [12] or [8,(3.33)]). It is not hard to show that Q is Lipschitz continuous,

$$|Q_{i+1/2}^n - q_i^n| \leq 2C_F |v_{i+1}^n - v_i^n|, \tag{6}$$

where C_F is the Lipschitz constant for F (cf. (4)). Assuming now that ϕ is non-negative, we obtain from the above discrete entropy inequality

$$\begin{aligned} -\Delta_T(v^{\Delta x}, \phi, k) &\leq -\Delta t \sum_{n=0}^N \sum_{i=-\infty}^{\infty} (Q_{i+1/2}^n - Q_{i-1/2}^n) \bar{\phi}_i^{n+1} + \Delta t \sum_{n=0}^N \sum_{i=-\infty}^{\infty} (q_{i+1}^n - q_i^n) \bar{\phi}_{i+1/2}^{n+1/2} \\ &= \Delta t \sum_{n=0}^N \sum_{i=-\infty}^{\infty} (Q_{i+1/2}^n - q_i^n) (\bar{\phi}_{i+1}^{n+1} - \bar{\phi}_i^{n+1}) \\ &\quad + \Delta t \sum_{n=0}^N \sum_{i=-\infty}^{\infty} (q_{i+1}^n - q_i^n) (\bar{\phi}_{i+1/2}^{n+1/2} - \bar{\phi}_{i+1}^{n+1}) \end{aligned}$$

(using (6) and the Lipschitz continuity $\|q\|_{\text{Lip}} \leq \|f\|_{\text{Lip}}$)

$$\leq \Delta t \sum_{n=0}^N \sum_{i=-\infty}^{\infty} |v_{i+1}^n - v_i^n| \left(C_F |\bar{\phi}_{i+1}^{n+1} - \bar{\phi}_i^{n+1}| + \|f\|_{\text{Lip}} |\bar{\phi}_{i+1/2}^{n+1/2} - \bar{\phi}_{i+1}^{n+1}| \right)$$

(smoothness of ϕ)

$$\leq \Delta t (C_F \Delta x \|\partial_x \phi\|_{L^\infty} + \|f\|_{\text{Lip}} \Delta t \|\partial_t \phi\|_{L^\infty}) \sum_{n=0}^N \sum_{i=-\infty}^{\infty} |v_{i+1}^n - v_i^n|.$$

From this estimate we obtain

$$\begin{aligned} -\Lambda_{\epsilon_0, \epsilon}(v^{\Delta x}, w) &= -\int_0^T \int_{\mathbb{R}} \Lambda_T(v^{\Delta x}, \omega_{\epsilon_0}(\cdot - s) \omega_\epsilon(\cdot - y), w) dy ds \\ &\leq \Delta t \int_0^T \int_{\mathbb{R}} (C_F \Delta x \|\omega_{\epsilon_0}(\cdot - s)\|_{L^\infty} \|\omega'_\epsilon(\cdot - y)\|_{L^\infty} \\ &\quad + \|f\|_{\text{Lip}} \Delta t \|\omega'_\epsilon(\cdot - s)\|_{L^\infty} \|\omega_\epsilon(\cdot - y)\|_{L^\infty}) \sum_{n=0}^N \sum_{i=-\infty}^{\infty} |v_{i+1}^n - v_i^n| dy ds \\ &\leq C \Delta t \left(\frac{C_F \Delta x}{\epsilon} + \frac{\|f\|_{\text{Lip}} \Delta t}{\epsilon_0} \right) \sum_{n=0}^N \sum_{i=-\infty}^{\infty} |v_{i+1}^n - v_i^n| \end{aligned}$$

for some constant $C > 0$ only depending on ω .

It remains to estimate $\nu(v^{\Delta x}, \epsilon_0)$. The standard estimate

$$|v_i^{n+1} - v_i^n| \leq \frac{\Delta t}{\Delta x} C_F (|v_{i+1}^n - v_i^n| + |v_i^n - v_{i-1}^n|)$$

yields

$$\begin{aligned} v(v^{\Delta x}, \epsilon_0) &\leq \Delta x \sum_i \max(\epsilon_0, \Delta t) \frac{1}{\Delta x} C_F (|v_{i+1}^n - v_i^n| + |v_i^n - v_{i-1}^n|) \\ &= 2C_F \max(\epsilon_0, \Delta t) \text{TV}(v^n). \end{aligned}$$

□

3 Convergence Rates for Irregular Data

With the Kuznetsov lemma and its corollary in place, we are now in place to prove convergence rates for (3) with irregular data. We start with some preliminaries in Sect. 3.1 before proving convergence rates in Sect. 3.

3.1 Preliminaries

We define the *discrete Lip⁺ (semi-)norm* as the sublinear functional

$$|v|_{\text{DLip}^+} := \sup_{i \in \mathbb{Z}} \frac{v_{i+1} - v_i}{\Delta x} \quad \text{for } v \in \ell^\infty(\mathbb{R}).$$

Following [13] (see also [14]), we say that a numerical flux function is (*strictly*) *Lip⁺ stable* if

$$|v^{n+1}|_{\text{DLip}^+} \leq \frac{1}{|v^n|_{\text{DLip}^+}^{-1} + \beta \Delta t} \tag{7}$$

for some $\beta \geq 0$ ($\beta > 0$, respectively) which is independent of $\Delta t, \Delta x$. Iterating (7), it holds in particular that

$$|v^n|_{\text{DLip}^+} \leq \frac{1}{|v^0|_{\text{DLip}^+}^{-1} + \beta t^n} \quad \forall n \in \mathbb{N}.$$

It was shown in [13] that the Lax–Friedrichs, Engquist–Osher and Godunov schemes are all strictly Lip⁺ stable. (The Roe scheme is non-strictly Lip⁺ stable.) The concept of Lip⁺ stability is motivated by the Oleinik entropy condition for conservation laws with strictly convex flux functions [15], which states that the Lip⁺ seminorm $|u|_{\text{Lip}^+} := \sup_{x \neq y} \frac{u(x) - u(y)}{x - y}$ of a solution of (1) should decrease over time at a rate proportional to t^{-1} ; more precisely,

$$|u(t)|_{\text{Lip}^+} \leq \frac{1}{|u_0|_{\text{Lip}^+}^{-1} + \beta_0 t}$$

where $0 \leq \beta_0 \leq f''(v)$ for all $v \in \mathbb{R}$.

For a function $g \in L^1(\mathbb{R})$ we define its total variation as

$$\text{TV}(g) = \sup \left\{ \int_{\mathbb{R}} g(x) \phi'(x) dx \mid \phi \in C_c^1(\mathbb{R}), \|\phi\|_{L^\infty} \leq 1 \right\}.$$

We say that a finite volume scheme is *total variation diminishing* (TVD) if for every $u_0 \in \text{BV}(\mathbb{R})$, we have $\text{TV}(v^{n+1}) \leq \text{TV}(v^n)$ for all $n \geq 0$. We say that the scheme is *monotone* if for all cell averaged initial data u^0, v^0 with $u_j^0 \leq v_j^0$ for all $j \in \mathbb{Z}$, we have $u_j^n \leq v_j^n$ for all $n > 0$ and $j \in \mathbb{Z}$.

Lemma 3 Let $u \in C_c^\alpha(\mathbb{R})$ for some $\alpha > 0$ and let $u^{\Delta x} \in \ell^1(\mathbb{Z})$ be the volume averages of u ,

$$u_i^{\Delta x} = \int_{C_i} u(x) dx \quad i \in \mathbb{Z}.$$

Then

$$\|u^{\Delta x} - u\|_{L^1(\mathbb{R})} \leq C \Delta x^\alpha, \tag{8a}$$

$$\text{TV}(u^{\Delta x}) \leq C \Delta x^{\alpha-1}, \tag{8b}$$

where C only depends on α and the size of the support of u .

Proof Let $K = \{i \in \mathbb{Z} \mid C_i \cap \text{supp } u \neq \emptyset\}$. Then

$$\begin{aligned} \|u^{\Delta x} - u\|_{L^1(\mathbb{R})} &= \sum_{i \in K} \int_{C_i} |u^{\Delta x}(x) - u(x)| dx = \sum_{i \in K} \int_{C_i} \left| \int_{C_i} u(y) - u(x) dy \right| dx \\ &\leq \sum_{i \in K} \int_{C_i} \int_{C_i} |u(y) - u(x)| dy dx \leq \|u\|_{C^\alpha} \sum_{i \in K} \int_{C_i} \int_{C_i} |x - y|^\alpha dy dx \\ &\leq \|u\|_{C^\alpha} \sum_{i \in K} \int_{C_i} \Delta x^\alpha dx \leq \|u\|_{C^\alpha} \sum_{i \in K} \Delta x^{\alpha+1} \\ &= C \Delta x^\alpha \end{aligned}$$

where $C = \|u\|_{C^\alpha} \Delta x |K|$ and $|K|$ is the Lebesgue measure of K . Similarly,

$$\text{TV}(u^{\Delta x}) = \sum_{i \in K} \left| \int_{C_i} u(x + \Delta x) - u(x) dx \right| \leq \sum_{i \in K} \|u\|_{C^\alpha} \Delta x^\alpha = C \Delta x^{\alpha-1}$$

for the same constant C as above. □

3.2 Convergence Rates

Without any assumptions on the scheme beyond being monotone, we can only prove convergence rates for initial data whose Hölder exponent is not smaller than $1/2$, which the following theorem makes precise.

Theorem 1 For a flux function $f \in C^1(\mathbb{R})$, let $u: \mathbb{R} \times [0, T] \rightarrow \mathbb{R}$ be the entropy solution of (1) with initial data $u_0 \in C_c^\alpha(\mathbb{R})$ for some $\alpha \in (0, 1)$. Let $(v_i^n)_{i,n}$ be generated by a monotone finite volume scheme (3) with initial data u_0 . Then

$$\|u(T) - v^{\Delta x}(T)\|_{L^1(\mathbb{R})} \leq C \Delta x^{\alpha-1/2} \tag{9}$$

for any $T > 0$, for some $C > 0$ only depending on f and u_0 .

Proof Since u_0 is Hölder continuous with exponent α , Lemma 3 implies that $\text{TV}(v_0^{\Delta x}) \leq C \Delta x^{\alpha-1} < \infty$, and since the scheme is TVD we get $\text{TV}(v^{\Delta x}(t^n)) \leq \text{TV}(v_0^{\Delta x})$. Hence,

$$\sum_{n=0}^N \text{TV}(v^{\Delta x}(t^n)) \Delta t \leq C T \Delta x^{\alpha-1}.$$

We note furthermore that

$$\|u_0 - v_0^{\Delta x}\|_{L^1(\mathbb{R})} \leq C \Delta x^\alpha.$$

CONVERGENCE RATES WITH UNBOUNDED TV




Fig. 1 Example evolution of fractional Brownian motion under Burgers’ equation with the Rusanov flux

Combining the above estimates with that of Lemma 2, we see that

$$\begin{aligned} \|u(T) - v^{\Delta x}(T)\|_{L^1(\mathbb{R})} &\leq 2C\Delta x^\alpha + C\Delta x^{\alpha-1}(2\epsilon + \epsilon_0\|f\|_{\text{Lip}} + 2C_F \max(\epsilon_0, \Delta t)) \\ &\quad + CT \left(\frac{C_F\Delta x}{\epsilon} + \frac{\|f\|_{\text{Lip}}\Delta t}{\epsilon_0} \right) \Delta x^{\alpha-1}. \end{aligned} \tag{10}$$

Choosing $\epsilon = \epsilon_0 = \Delta x^{1/2}$ yields

$$\|u(T) - v^{\Delta x}(T)\|_{L^1(\mathbb{R})} \leq 2C_1\Delta x^\alpha + C_2\Delta x^{\alpha-1/2} \leq C\Delta x^{\alpha-1/2}, \tag{11}$$

which was what we wanted. \square

We can improve the somewhat suboptimal convergence rate of $\Delta x^{\alpha-1/2}$ for Lip^+ -stable schemes. To this end we need the following result.

Lemma 4 *Let $u_0 \in L^1 \cap L^\infty(\mathbb{R})$ have compact support and let u be the entropy solution of (1). Let $(v^n)_{i,n}$ be generated by a strictly Lip^+ stable finite volume scheme (3). Then*

$$\sum_{n=0}^N \text{TV}(v^n)\Delta t \leq C \left(|v^0|_{\text{DLip}^+} \Delta t + \frac{1}{\beta} \log(1 + \beta t^N |v^0|_{\text{DLip}^+}) \right)$$

where $t^N \leq T$ and $C > 0$ is independent of Δx .

Proof Let $M > 0$ be such that $\text{supp } v^{\Delta x}(t) \in [-M, M]$ for all $t \in [0, T]$ and let $I \in \mathbb{N}$ be such that $x_{I-1} < M \leq x_I$. The compact support of v and the strict Lip^+ stability imply that

$$\text{TV}(v^n) = \sum_{i=-I}^I |v_{i+1}^n - v_i^n| = 2 \sum_{i=-I}^I (v_{i+1}^n - v_i^n)^+ \leq \sum_{i=-I}^I \frac{1}{|v^0|_{\text{DLip}^+}^{-1} + \beta t^n} \Delta x \leq 2M \frac{1}{|v^0|_{\text{DLip}^+}^{-1} + \beta t^n}$$

U. S. FJORDHOLM AND K. O. LYE




Fig. 2 Comparison between different mesh resolutions ($\Delta x = 2^{-15}$ and $\Delta x = 2^{-8}$) at $T = 0$ and $T = 1$ for different equations and schemes. Here $H = 0.125$. In the left column, we show the initial data, while in the right column, we show the evolved data under the specified numerical scheme

Hence,

$$\begin{aligned}
 \sum_{n=0}^N \text{TV}(v^n) \Delta t &\leq 2M \sum_{n=0}^N \frac{1}{|v^0|_{\text{DLip}^+}^{-1} + \beta t^n} \Delta t \\
 &\leq 2M \left(\frac{\Delta t}{|v^0|_{\text{DLip}^+}^{-1}} + \frac{1}{\beta} \left(\log(|v^0|_{\text{DLip}^+}^{-1} + \beta t^N) - \log(|v^0|_{\text{DLip}^+}^{-1}) \right) \right) \\
 &= 2M \left(|v^0|_{\text{DLip}^+} \Delta t + \frac{1}{\beta} \log(1 + \beta t^N |v^0|_{\text{DLip}^+}) \right). \tag{12}
 \end{aligned}$$





Fig. 3 Comparison between different mesh resolutions ($\Delta x = 2^{-15}$ and $\Delta x = 2^{-8}$) at $T = 0$ and $T = 1$ for different equations and schemes. Here $H = 0.5$. In the left column, we show the initial data, while in the right column, we show the evolved data under the specified numerical scheme

□


Theorem 2 For a strictly convex flux function $f \in C^1(\mathbb{R})$, let $u : \mathbb{R} \times [0, T] \rightarrow \mathbb{R}$ be the entropy solution of (1) with initial data $u_0 \in C_c^\alpha(\mathbb{R})$ for some $\alpha \in (0, 1)$. Let $(v_i^n)_{i,n}$ be generated by a strictly Lip^+ stable, monotone finite volume scheme (3) with initial data u_0 . Then

$$\|u(T) - v^{\Delta x}(T)\|_{L^1(\mathbb{R})} \leq C_{L,M,f,\beta} \sqrt{\log(1 + C_F \beta T \Delta x^{\alpha-1})} \Delta x^{\alpha/2} \tag{13}$$


U. S. FJORDHOLM AND K. O. LYE




(A) Burgers with Godunov



(B) Burgers with Rusanov



(C) Cubic with Rusanov



(D) Linear with Rusanov

Fig. 4 Convergence rates for different equations and numerical fluxes at $T = 1$

U. S. FJORDHOLM AND K. O. LYE




Fig. 5 The total variation as a function of time for varying equations, numerical fluxes and Hurst indices. This corresponds to the first estimate in (12)

for any $T > 0$, for some $C_{L,M,f,\beta} > 0$. For small $\Delta x > 0$ this yields the “almost $\Delta x^{\alpha/2}$ ” estimate

$$\|u(T) - v^{\Delta x}(T)\|_{L^1(\mathbb{R})} \leq C \Delta x^{\alpha/2} \sqrt{-\log \Delta x}. \tag{14}$$

Proof Lemmas 4 and 3 imply

$$\sum_{n=0}^N \text{TV}(v^n) \Delta t \leq C \left(\|u_0\|_{C^\alpha} \Delta x^\alpha + \frac{1}{\beta} \log(1 + \|u_0\|_{C^\alpha} \beta t^N \Delta x^{\alpha-1}) \right)$$

for some $C > 0$ independent of Δx . Inserting this and the bounds $\|u_0 - v_0^{\Delta x}\|_{L^1(\mathbb{R})} \leq C \Delta x^\alpha$ and $\text{TV}(v_0^{\Delta x}) \leq C \Delta x^{\alpha-1}$ from Lemma 3 into the Kuznetsov estimate (5), we produce

$$\begin{aligned} \|u(T) - v^{\Delta x}(T)\|_{L^1(\mathbb{R})} &\leq C \Delta x^\alpha + C \Delta x^{\alpha-1} (\epsilon + \epsilon_0 \|f\|_{\text{Lip}} + \max(\epsilon_0, \Delta t)) \\ &+ C \left(\frac{C_F \Delta x}{\epsilon} + \frac{\|f\|_{\text{Lip}} \Delta t}{\epsilon_0} \right) \left(\Delta x^\alpha + \frac{1}{\beta} \log(1 + \|u_0\|_{C^\alpha} \beta T \Delta x^{\alpha-1}) \right). \end{aligned} \tag{15}$$

Defining

$$Q(\Delta x, \alpha) = \Delta x^\alpha + \frac{1}{\beta} \log(1 + \|u_0\|_{C^\alpha} \beta T \Delta x^{\alpha-1})$$




Fig. 6 Sharpness of the bound (12) for Burgers’ equation solved with the Godunov scheme. Specifically, we plot the ratio $\frac{2M(|v^0|_{DLip} + \Delta t + \frac{1}{\beta} \log(1 + \beta t^N |v^0|_{DLip}))}{\sum_{n=0}^N TV(v^n) \Delta t}$ as a function of the mesh width Δx . A value of 1 corresponds to a sharp estimate, a value greater than one corresponds to an not sharp estimate

and setting $\epsilon = \epsilon_0 = \Delta x^{1-\alpha/2} \sqrt{Q(\Delta x, \alpha)}$ yields

$$\|u(T) - v^{\Delta x}(T)\|_{L^1(\mathbb{R})} \leq C \Delta x^\alpha + C_{L,M,f} \sqrt{Q(\Delta x, \alpha)} \Delta x^{\alpha/2} \tag{16}$$

Since $\alpha \in (0, 1)$, the second term on the right-hand side dominates, so we obtain (13). To get (14) we estimate

$$\log(1 + C \Delta x^{\alpha-1}) \lesssim \log(C) + \log(\Delta x^{\alpha-1}) \lesssim -(1 - \alpha) \log(\Delta x).$$

□

4 Numerical Examples

We consider three scalar conservation laws: Burgers’ equation where $f(u) = u^2/2$, a cubic conservation law where $f(u) = u^3/3$, and lastly a linear conservation law where $f(u) = u$. The initial data will be given as fractional Brownian motion with varying Hurst exponent H . Introduced by Mandelbrot et al. [16], fractional Brownian motion can be seen as a generalization of standard Brownian motion with a scaling exponent different than 1/2. We set

$$u_0^H(x) := B^H(x) \quad x \in [0, 1],$$

where B^H is a path of fractional Brownian motion with Hurst exponent $H \in (0, 1)$. Brownian motion corresponds to a Hurst exponent of $H = 1/2$.

To generate fractional Brownian motion, we use the random midpoint displacement method originally introduced by Lévy [17] for Brownian motion, and later adapted for fractional Brownian motion [18, 19]. For a more detailed description on how we generate fractional Brownian motion, consult [20,A.4.1]. See Fig. 1 (left column) for an example

with $H = 0.125$, $H = 0.5$ and $H = 0.75$. The initial data is normalized to have values in $[-1, 1]$.

4.1 Numerical Results

Figures 2 and 3 show the computed solutions for mesh resolutions of $\Delta x = 2^{-8}$ and $\Delta x = 2^{-15}$, and as expected the approximation converges upon mesh refinement. In order to measure the rate of convergence we compare with a reference solution computed on a mesh of 2^{16} cells ($\Delta x = 2^{-16}$); the results are shown in Fig. 4. The figure clearly shows convergence for all the given configurations. However, for most configurations—most notably those of low Hurst index H —we observe better convergence rates than the rates $\Delta x^{H-1/2}$ and $\Delta x^{H/2} \sqrt{|\log \Delta x|}$ predicted by Theorems 12, respectively.

4.2 Sharpness of our Estimates

In Fig. 5 we show the evaluation of the inverse of the total variation as a function of time. As we can see from the plot, the total variation decays as C/t , which agrees well with the estimate on $\text{TV}(v^n)$ in the proof of Lemma 4.

Inspired by (12), we use the value of $\beta = \frac{1}{2} \cdot \frac{1}{4}$ for the Godunov flux applied to Burgers' equation, computed in [13], and measure the sharpness of the bound (12) by the ratio

$$\frac{2M \left(|v^0|_{\text{DLip}^+} \Delta t + \frac{1}{\beta} \log \left(1 + \beta t^N |v^0|_{\text{DLip}^+} \right) \right)}{\sum_{n=0}^N \text{TV}(v^n) \Delta t},$$

which is plotted in Fig. 6 as a function of the spatial resolution Δx . As we can see, the bound is not perfectly sharp, and seems to overestimate the sum on the left hand side of (12) by a factor of $\Delta^{\frac{1}{4}}$. This partially explains the discrepancy of the observed convergence rates and the predicted convergence rate of Theorem 2.

5 Conclusion

In this paper we have extended the standard Kuznetsov convergence proof for finite volume schemes approximating solutions to hyperbolic conservation laws to include initial data with unbounded total variation. The theory covers rapidly oscillating data such as Brownian and fractional Brownian motion. We show several numerical experiments which show good agreement with the theory.

Our result can easily be extended to cover initial data which is only piecewise Hölder continuous with a finite number of downward jump discontinuities. The suboptimal rate $\Delta x^{\alpha-1/2}$ in Theorem 1 can be extended even further to cover e.g. initial data in Besov spaces, since it only relies on the projection estimates (8a) and (8b).

We conjecture that the rate $\Delta x^{\alpha/2}$ given in (14) in Theorem 2 is optimal or near optimal. For optimality of convergence rates see e.g. [21, 22].

Funding Open access funding provided by University of Oslo. USF was partially supported by the Research Council of Norway project number 301538.

Data and Code Availability All numerical experiments have been done with the Alsvinn code available from <https://alsvinn.github.io>. The code for post-processing, along with the generation of the initial data, can be

found at https://github.com/kjetil-lye/unbounded_tv_experiments. They are also permanently stored on the Zenodo platform with the DOI 10.5281/zenodo.4088164.

Declarations

Conflict of interest The Author declare that they have no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Coclite, G.M., Mishra, S., Risebro, N.H., Weber, F.: Analysis and numerical approximation of brinkman regularization of two-phase flows in porous media. *Comput. Geosci.* **18**(5), 637–659 (2014)
2. Krotkiewski, Marcin, Ligaarden, Ingeborg S., Lie, Knut-Andreas., Schmid, Daniel W.: On the importance of the stokes-brinkman equations for computing effective permeability in karst reservoirs. *Commun. Comput. Phys.* **10**(5), 1315–1332 (2011)
3. Sinai, Y.G.: Statistics of shocks in solutions of inviscid burgers equation. *Commun. Math. Phys.* **148**(3), 601–621 (1992)
4. Bec, J., Khanin, K.: Burgers turbulence. *Phys. Rep.* **447**, 1–66 (2007)
5. She, Z.-S., Aurell, E., Frisch, U.: The inviscid Burgers equation with initial data of Brownian type. *Commun. Math. Phys.* **148**, 623–641 (1992)
6. Weber, F.: Convergence rates of finite difference schemes for the linear advection and wave equation with rough coefficient. *IMA J. Numer. Anal.* **37**(3), 1586–1634 (2016)
7. Dafermos, C.M.: Hyperbolic conservation laws in continuum physics. In: *Fundamental Principles of Mathematical Sciences*, Vol. 325, 4th ed. Springer-Verlag, Berlin (2016)
8. Holden, H., Risebro, N.H.: Front tracking for hyperbolic conservation laws. In: *Applied Mathematical Sciences*. Vol 152, 2nd ed. Springer, Heidelberg (2015)
9. LeVeque, R.J.: Numerical methods for conservation laws. *Lectures in Mathematics ETH Zürich*, 2nd ed. Birkhäuser Verlag, Basel (1992)
10. Kuznetsov, N.N.: Accuracy of some approximate methods for computing the weak solutions of a first-order quasi-linear equation. *USSR Comput. Math. Math. Phys.* **16**(6), 105–119 (1976)
11. Kruzkov, S.N.: First order quasilinear equations in several independent variables. *Math USSR SB* **10**(2), 217–243 (1970)
12. Crandall, M.G., Majda, A.: Monotone difference approximations for scalar conservation laws. *Math. Comput.* **34**, 1–21 (1980)
13. Nessyahu, H., Tadmor, E.: The convergence rate of approximate solutions for nonlinear scalar conservation laws. *SIAM J. Numer. Anal.* **29**(6), 1505–1519 (1992)
14. Fjordholm, U.S., Solem, S.: Second-order convergence of monotone schemes for conservation laws. *SIAM J. Numer. Anal.* **54**(3), 1920–1945 (2016)
15. Oleinik, O.A.: On discontinuous solutions of non-linear differential equations. *Uspekhi Mat. Nauk.* **12**:3–73, 1957
16. Mandelbrot, Benoit B., Van Ness, John W.: Fractional Brownian motions, fractional noises and applications. *SIAM Rev.* **10**, 422–437 (1968)
17. Lévy, P.: *Processus stochastiques et mouvement brownien*. Les Grands Classiques Gauthier-Villars. Éditions Jacques Gabay, Sceaux (1992)
18. Fournier, A., Fussell, D., Carpenter, L.: Computer rendering of stochastic models. *Commun. ACM* **25**(6), 371–384 (1982)
19. Voss, R.F.: Random fractal forgeries. In: Earnshaw, R.A. (ed.) *Fundamental Algorithms for Computer Graphics* NATO Advanced Study Institute Directed by J.E. Bresenham, R.A. Earnshaw, M.L.V. Pitteway, pp. 805–835. Springer Berlin Heidelberg, Berlin (1991)

20. Lye, K.O.: Computation of statistical solutions of hyperbolic systems of conservation laws. PhD thesis, ETH Zürich (2020)
21. Şabac, F.: The optimal convergence rate of monotone finite difference methods for hyperbolic conservation laws. *SIAM J. Numer. Anal.* **34**(6), 2306–2318 (1997)
22. Ruf, A.M., Sande, E., Solem, S.: The optimal convergence rate of monotone schemes for conservation laws in the Wasserstein distance. *J. Sci. Comput.* **80**(3), 1764–1776 (2019)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.