

1 **Estimating overall size-selection pattern in the bottom trawl**
2 **fishery for four economically important fish species in the**
3 **Mediterranean Sea**

4
5 Chryssi Mytilineou^{1,2*}, Bent Herrmann³, Antonello Sala⁴, Danai Mantopoulou-Palouka¹ and
6 Persefoni Megalofonou²

7
8 *¹Hellenic Centre for Marine Research (HCMR), Institute of Marine Biological Resources &*
9 *Inland Waters (IMBRIW), P.O. Box 712, Anavyssos 19013, Attica, Greece*

10 *²Faculty of Biology, Department of Zoology-Marine Biology, National and Kapodistrian*
11 *University of Athens (NKUA), Panepistimioupolis, Ilissia, 15784 Athens, Greece*

12 *³SINTEF Ocean, Fishery Technology, Willemoesvej 2, 9850 Hirtshals, Denmark; University*
13 *of Tromsø, Norway; DTU Aqua, Technical University of Denmark, Hirtshals, Denmark*

14 *⁴Italian National Research Council (CNR), Institute of Marine Biological Resources and*
15 *Biotechnologies (IRBIM), Italy*

16

17

18

19

20

21

22

23 ***Corresponding author:**

24 Mytilineou Chryssi

25 *E-mail:* chryssi@hcmr.gr

26 *tel.:* +30. 2111065244

27 *fax:* +30. 2111065256

28 Postal address: Hellenic Centre for Marine Research, Institute of Marine Biological
29 Resources & Inland Waters, 46.7 km Athens-Sounio Av., P.O. Box 712, Anavyssos 19013,
30 Attica, Greece

31 ORCID ID: <https://orcid.org/0000-0002-9326-1650>

32

33 **ABSTRACT**

34 The management of multispecies fisheries, such as the Mediterranean bottom trawl fishery, is
35 always a challenge. However, information on gear selectivity and discards has been studied
36 separately so far. In this paper, the overall size-selection pattern by the trawl codend in the
37 sea and by the fisher onboard the vessel is investigated for four commercially important fish
38 species, *Mullus barbatus*, *Mullus surmuletus*, *Pagellus erythrinus* and *Lophius budegassa*,
39 using different codends. For each species, the selection model used offered the possibility to
40 simultaneously describe the escape, discard, and landing probability. The results, useful for
41 fisheries management, showed that the codend made of 40 mm diamond meshes was always
42 detrimental for the stocks. The 40 mm square meshes codend compared to that of 50 mm
43 diamond meshes was more appropriate for the sustainability of both *Mullus* species,
44 providing also a lawful catch along with greater compliance to the rules fisher behaviour,
45 negligible discards and the lowest possible economic losses for the fisher. None of the
46 codends was effective for *P. erythrinus* in achieving the minimum conservation reference size
47 (MCRS) of the species. All codends were harmful to *L. budegassa* as the majority of
48 juveniles were retained in the codend, resulting in negligible escapees, a high discard
49 probability, and landings of a size much lower than the length at first maturity of the species.
50 Further studies are needed to be conducted in the future for other species, since the trawl
51 fishery in the Mediterranean is a multi-species fishery.

52

53

54

55 **Keywords:**

56 size selection, discards, mullet, common pandora, blackbellied anglerfish

57

58

59 **Highlights**

- 60 • **Gear and fisher selection processes may be described by an overall selection model**
- 61 • **Selection models can predict escape, discard and landing probability concurrently**
- 62 • **40mm square mesh in Mediterranean trawl codend is more sustainable for stocks**
- 63 • **40mm square meshes trawl codend produces negligible red/stripped mullet**
- 64 **discards**
- 65 • **Anglerfish sustainability cannot be ensured by the current Common Fishery**
- 66 **Policy**

67 1. Introduction

68 The Mediterranean trawl fishery primarily targets species of high economic importance,
69 such as hake (*Merluccius merluccius*), mullets (*Mullus* spp.), common pandora (*Pagellus*
70 *erythrinus*) and blackbellied anglerfish (*Lophius budegassa*). Bycatch of juveniles of these
71 species ends up discarded (Tsagarakis et al. 2017; Bellido et al. 2017; Mytilineou et al. 2018,
72 2020). Juvenile protection and the reduction of undersized catch below the minimum
73 conservation reference size (MCRS) are important issues in the European Common Fishery
74 Policy, particularly related to the management of the Mediterranean bottom trawl fishery
75 (Council Regulation (EC) No 1967/2006; Regulation (EU) 2019/1241). Gear and fisher
76 selection patterns are related to these issues affecting stock sustainability (Vasilakopoulos et
77 al. 2015). Although gear selectivity and discard probability have generally been studied
78 separately, Mytilineou et al. (2018) combined these two sequential selection processes and
79 modelled for the first time the overall selection process on a fish population entering the trawl
80 codend. This approach first applied for European hake, Atlantic horse mackerel and four-
81 spotted megrim, and based on selectivity data, simultaneously predicts the escape, discard,
82 and landing probability of the species studied. As the Mediterranean trawl fishery is a
83 multispecies one, such information is essential for all target species, especially under the state
84 of overexploitation of most stocks in the Mediterranean (FAO, 2020).

85 In the Mediterranean, many studies have been conducted for the trawl codend selectivity
86 of red mullet (e.g. Tokaç et al. 2014; Sala et al. 2015) and common pandora (e.g. Ateş et al.
87 2010; Özbilgin et al. 2012); only one for striped mullet (Ordines et al. 2006) and none for
88 blackbellied anglerfish. On the other hand, several studies on the discard probability of these
89 species have been conducted, based on data from observers onboard fishing vessels (e.g.
90 Tsagarakis et al. 2017; Damalas et al. 2018). To date, no research has provided combined
91 information on the overall selection of these species for the Mediterranean trawl fishery.

92 The objective of this study is twofold: i) to investigate the potential applicability of the
93 model proposed by Mytilineou et al. (2018) based on the population of red mullet (*Mullus*
94 *barbatus*), striped mullet (*Mullus surmuletus*), common pandora (*Pagellus erythrinus*) and
95 blackbellied anglerfish (*Lophius budegassa*) entering the trawl codend and ii) to study the
96 overall size-selection for these commercially important species and provide information on
97 their escape, discard and landing probability.

98

99 **2. Materials and methods**

100 *2.1. Data collection*

101 From September 9 to October 4, 2014, a selectivity experimental survey was conducted on
102 fishing grounds of the South Aegean Sea (Fig. 1, for details Table S1 in Supplementary
103 material). For this purpose, a commercial trawler, equipped with a bottom trawl used in
104 professional fishing, was hired. The specifications of the gear are described in detail in the
105 Supplementary material as well as in Mytilineou et al. (2018). The depth range of the
106 experimental fishing was between 50 and 310 m, in line with the main depth range in which
107 the commercial Greek trawl fleet operates. Fishing was carried out in 28 locations using three
108 different codends resulting in a total of 84 hauls (3 X 28). Invalid hauls due to damaged net
109 or poor gear performance during fishing, which was checked by SCANMAR, were excluded
110 from the analysis.

111 Three codends were used to conduct the experimental fishing survey: i) a codend of 40
112 mm nominal size square meshes (40S), which has been established in the commercial
113 Mediterranean trawl fishery according to the Council Regulation (EC) No 1967/2006 (actual
114 mesh size: 43.2 ± 0.6 mm), (ii) a codend of 50 mm nominal size diamond meshes (50D),
115 which can be used in accordance to the above-mentioned regulation, if it is more selective
116 than the 40S (actual mesh size: 51.1 ± 0.7 mm) and (iii) a codend of 40 mm nominal size

117 diamond meshes (40D) (actual mesh size: 43.2 ± 0.6 mm); the latter, although prohibited in
118 EU Mediterranean countries, was investigated because smaller or slightly larger meshes are
119 still in use in various Mediterranean trawl fleets (e.g. Ragheb et al. 2019). In all cases, the
120 three codends were 5.6 m in length, and with the same circumferential length at sea (~4.3 m).
121 They were knotless, and made by single twine multifilament nylon (PA) of 2.8 mm twine
122 thickness. The number of meshes in codend circumference was 400, 200, and 340 meshes for
123 the 40D, the 40S, and 50D, respectively. The characteristics of the three codends and their
124 meshes are described in detail in the Supplementary material (Table S2).

125 Data were collected for four species, red mullet *M. barbatus* Linnaeus, 1758, striped mullet
126 *M. surmuletus* Linnaeus, 1758, common pandora *P. erythrinus* (Linnaeus, 1758) and
127 blackbellied anglerfish *L. budegassa* Spinola, 1807, which were selected for their commercial
128 importance and high economic value, and therefore the need of information for their
129 sustainable exploitation and management. *M. barbatus*, *M. surmuletus*, *P. erythrinus* and *L.*
130 *budegassa* are species with different body shape characteristics; the first two of rounded body
131 shape, the third one very compressed and of high body depth, and the last one with a very
132 large head. Furthermore, the first three species are regulated with MCRS at 11 cm, 11 cm and
133 15 cm, respectively (Council Regulation (EC) No 1967/2006; Regulation (EU) 2019/1241). *L.*
134 *budegassa*, although regulated in the past with MCRS at 30 cm TL (Council Regulation (EC)
135 No 1626/94), is no longer part of any new regulation nowadays. Moreover, a policy reform of
136 the legislated discard ban, permitted *M. barbatus* discarding up to 6% of the total annual
137 landings of the species by 2019 (Commission Delegated Regulation (EU) 2017/86). Apart
138 from the MCRS, the length at first maturity of these species, available in the literature, was
139 also used as a threshold for the sustainable exploitation of their stocks by the trawl.

140 A three-compartment sampling scheme was used to classify escapees, discards, and
141 landings as described in Mytilineou et al. (2018). The cover-codend method (Wileman *et al.*,

142 1996) was used to collect the data for the escapees. The design for the cover was similar to
143 that presented by Sala *et al.* (2015) using in the cover a 20-mm diamond mesh size net.
144 During the process, escapees were retained in the cover, while the separation between
145 discards and landings as well as the classification of the landings compartment into different
146 commercial categories was performed by the vessel crew simulating the procedure followed
147 in commercial fishing. The landings compartment, depending on the species, was divided into
148 two or three compartments according to their size related commercial value (i.e category A, B,
149 and C) for further analyses.

150

151 2.2. Predicting the overall size-selection process

152 The methodological approach for modelling the sequential size-selection processes, both in
153 the sea and onboard the fishing vessel, is described in detail in Mytilineou *et al.* (2018). In
154 summary, a fish of length l after entering the gear in the sea has a probability p_{esc} to escape
155 through the codend, or equivalently:

$$156 \quad p_{esc}(l) = 1 - r_{gear}(l)$$

157 where r_{gear} is the probability of a fish to be retained by the gear. Then, given that the fish is
158 retained, it has a probability p_{land} to be landed by the fisher. Denoting by r_{fisher} , the probability
159 of a retained by the gear fish being retained by the fisher and landed, we have:

$$160 \quad p_{land}(l) = r_{gear}(l) \times r_{fisher}(l)$$

161 We then denote by p_{disc} the probability of a fish to be discarded by the fisher, given that it has
162 been retained by the gear. The mathematical formulation of this process can be described as
163 follows:

$$164 \quad p_{disc}(l) = (1 - r_{fisher}(l)) \times r_{gear}(l)$$

165 Since both probabilities, p_{esc} and p_{land} can be interpreted as size selection procedures and
166 given that in most cases smaller fish are being discarded, their probabilities are represented
167 by sigmoid curves, while the p_{disc} is fitted by a bell-shaped curve.

168 Following Wileman et al. (1996), selection curves can be adequately described by four
169 different models: Logit, Probit, Gombertz and Richard. In the present analysis, the four
170 models were fitted to the data of each sequential selection process r_{gear} and r_{fisher} . A total of
171 16 different combinations of models were tested for each codend. The best model was
172 selected based on the p-value (should be >0.05) as well as the model deviance compared to
173 the degrees of freedom (Wileman et al., 1996), followed by the AIC criterion (Akaike, 1974).
174 These models are described by a set of parameters: the length at which 50% of the fish is
175 being retained either by the codend or the fisher, denoted as $L50_{gear}$ and $L50_{fisher}$ respectively,
176 the selection range SR_{gear} and SR_{fisher} (denoted as the difference $L75 - L25$) and in the case of
177 Richard model an additional parameter δ , which describes the asymmetry of the curve. Let ν
178 denote the set of parameters for each model. Then the probabilities of r_{gear} and r_{fisher} can be
179 expressed as:

$$180 \quad r_{gear}(l, \nu_{gear}) \text{ and } r_{fisher}(l, \nu_{fisher})$$

181 Since the probability p_{land} is expressed by the two curves r_{gear} and r_{fisher} , the parameters
182 $L50_{land}$ and SR_{land} can also be estimated. This method was described in Sistiaga et al. (2010).
183 Parameter estimation for the three different probabilities: p_{esc} , p_{disc} and p_{land} was performed
184 using of the maximum log-likelihood function as applied by Mytilineou et al. (2018).
185 Although a mean selection curve is generally estimated on individual haul basis (Fryer,
186 1991), in the present study average selection parameters were estimated for each codend by
187 pooling the data of the hauls, as proposed by Millar (1993) for fisheries. However here, the
188 three compartments design was considered for the two sequential selection processes of the
189 overall selection model (see equation described in Mytilineou et al., 2018). Besides the

190 average selectivity curve, a bootstrap technique was applied to calculate the "Efron percentile
191 95% confidence limits" (95% CI) for this curve (Efron, 1982), taking into account both
192 within and between haul variation (Millar, 1993). All the analysis described above was
193 implemented using SELNET software (Hermann et al. 2012, 2013) and applied in several
194 works (Sala et al. 2015; Mytilineou et al. 2018; Herrmann et al. 2019).

195

196 2.3. Comparisons between gears

197 The parameters of the three codends were compared through the overlapping of their
198 estimated 95% CIs (Frandsen et al., 2010). Furthermore, length-dependent differences
199 between the three codends were calculated for the probabilities p_{esc} , p_{disc} , and p_{land} using the
200 following formulas:

$$201 \quad \Delta p_{esc}(l) = p_{esc_i}(l) - p_{esc_j}(l)$$

$$202 \quad \Delta p_{disc}(l) = p_{disc_i}(l) - p_{disc_j}(l)$$

$$203 \quad \Delta p_{land}(l) = p_{land_i}(l) - p_{land_j}(l)$$

204 where l is the length class and $j, i = (40D, 40S, 50D)$ with $i \neq j$. The differences were
205 accompanied by their related Efron 95% confidence limits. In the case that the 95% CI of a
206 length class includes zero, the difference is not statistically significant. The method was
207 applied by several researchers (Larsen et al. 2019; Mytilineou et al. 2020).

208

209 2.4. Discard Indicators

210 Discard indicators proposed by Mytilineou et al. (2018) were also estimated in this work.
211 Specifically: LDp_{max} is the length at which the probability of a fish to be discarded is the
212 highest, denoted as Dp_{max} ; $DR_{0.05}$, $DR_{0.25}$, $DR_{0.50}$, $DR_{0.75}$, $DR_{0.95}$ are the different ranges of the
213 discard bell-shaped curve at different values of probability and $DA_{0.05}$ is the surface of the

214 discard bell-shaped curve when the probability is ≥ 0.05 (for details see Fig. 3 in Mytilineou
215 et al. 2018).

216

217 **3. Results**

218 *3.1. Experimental data*

219 Data for *M. barbatus*, *M. surmuletus*, *P. erythrinus* and *L. budegassa* collected during the
220 experimental survey per haul and in total for the three compartments, the escapees, the
221 discards, and the landings for each codend and their percentage to the total amount of the
222 species entering the trawl codend are presented in Tables S3-S6 in the Supplementary
223 material Both *M. barbatus* and *M. surmuletus* were caught in many hauls (18 and 12,
224 respectively) and in high numbers (Table S3 and S4). *P. erythrinus*, although being fished in 8
225 of the hauls, was also collected in high numbers (Table S5). *L. budegassa* was caught in 9 of
226 the hauls, but in very low numbers, reflecting the generally low abundance of the species in
227 the sea (Table S6).

228

229 *3.2. Mullus barbatus*

230 The parameters and the statistics for the best overall selection model of *M. barbatus*
231 appear in Table 1. The model fitted the data well in all cases (Fig. 2).

232 The $L50_{gear}$ of *M. barbatus* was significantly higher for the 40S than for the two diamond
233 codends. Similar $L50_{gear}$ with overlap of their 95% CI was found for the 40D and 50D
234 (Table 1). Significantly higher escape probability was detected for the 40S and 50D compared
235 to 40D mainly for lengths ranging between 9 - 17 cm and 11 - 23 cm TL, respectively (Fig.
236 3). The escape probability of the 50D compared to 40S was significantly lower for lengths 9 -
237 13 cm, but significantly higher for lengths ≥ 17 cm TL (Fig. 3), related to the higher SR_{gear}
238 value of the 50D (Table 1).

239 The discard probability showed relatively low values in all cases (Fig. 2), indicating that a
240 few *M. barbatus* entering the three codends will be discarded. Statistically significant higher
241 discard probability for the 40D and 50D compared to the 40S was predicted for the sizes ≥ 12
242 cm TL; however, of negligible importance (Fig. 3). The 40S codend showed the lowest
243 discard indicators with Dp_{max} at 0.06 and $DA_{0.05}$ almost zero, but an overlap of the 95% CI
244 was detected for all of them among the three codends (Table 2).

245 $L50_{land}$ of *M. barbatus* was significantly higher for the 40S and lower for the 40D (Table
246 1); an overlap of the 95% CI of the latter was found with that of 50D. Fisher landing
247 probability displayed significantly higher values for the 40D than the 40S at lengths 9 - 18 cm
248 TL (most important between 9 – 15 cm); higher for the 40D than 50D at lengths ≥ 10 cm TL
249 (most important between 10 – 20 cm). Landing probability was also higher for the 50D than
250 40S for the sizes 10 – 13 cm TL, but higher for the 40S than 50D at sizes ≥ 17 cm TL (Fig. 2,
251 3).

252

253 3.3. *Mullus surmuletus*

254 The overall selection model fitted the data of *M. surmuletus* well, although the very small
255 number of individuals and the absence of some size classes in the 40D escapees and discards
256 produced large 95% CI (Fig. 2). The results for the best model appear in Table 1.

257 $L50_{gear}$ of *M. surmuletus* was significantly higher for the 40S than the other two codends.
258 Overlap of their 95% CI was found for the 40D and 50D (Table 1). A significantly higher
259 escape probability was detected for the 40S and the 50D compared to 40D, mainly for lengths
260 between 8 - 15 cm and 10 - 25 cm TL, respectively (Fig. 4). The escape probability of the
261 50D codend compared to 40S was significantly lower for lengths 8 - 11 cm, but significantly
262 higher especially for lengths 15 - 25 cm TL (Fig. 4), which is related to the higher SR_{gear}
263 value of the 50D (Table 1).

264 The discard probability of *M. surmuletus* was significantly higher for the two diamond
265 codend than the 40S one; the latter without discards (Fig. 2). Statistically significant higher
266 discard probability for the 40D and 50D compared to 40S was predicted for the sizes 5 - 11
267 cm TL; no significant difference between the diamond codends (Fig. 4). Similar indicators
268 were obvious for the 40D and 50D, which differed significantly from the zero values of 40S
269 (Table 2).

270 The parameter $L50_{land}$ was significantly higher for the 40S and lower for the 40D (Table
271 1). An overlap of the 95% CI of the latter was observed with that of the 50D. Fisher landing
272 selection displayed significantly higher values for the 40D than the 40S or 50D at lengths >10
273 cm (most important between 10 - 15 cm). It was higher for the 50D than 40S only for the size
274 11 cm TL, but higher for the 40S than 50D at sizes ≥ 15 cm (most important between 15 - 25
275 cm) (Fig. 4).

276

277 3.4. *Pagellus erythrinus*

278 The parameters and the statistics for the best overall selection model of *P. erythrinus*
279 appear in Table 3. The model generally fitted the data well, although large 95% CI were
280 obtained for the 40D escape and landing probability as a result of the very low number of
281 individuals in these cases (Fig. 5).

282 The gear selection parameter of *P. erythrinus* was higher for the 50D codend, although
283 overlap of the 95% CI was obvious between the 40D and 40S and between the 40S and 50D
284 (Table 3). A significantly higher escape probability was detected for the 40S and 50D
285 compared to the 40D at lengths >10 cm (mainly between 10 - 17 cm) (Fig. 6). The escape
286 probability of the 50D codend compared to 40S was significantly higher at lengths between
287 13 - 16 cm TL (Fig. 6), associated with the higher SR_{gear} of the former codend (Table 3).

288 The discard probability of *P. erythrinus* showed higher values for the 40D codend (Fig. 5).
289 Significantly higher values for the two diamond codends compared to the 40S were predicted
290 for sizes ≤ 5 cm (Fig. 6). Comparison between the 40D and 50D revealed statistically
291 significant higher discard probability for the sizes 10 - 11 cm TL for the former codend (Fig.
292 6). Some of the discard indicators of the 50D were lower than those of the other two codends,
293 however, overlap in their 95% CI was obvious in all cases among the three codends (Table 2).

294 The landing probability of *P. erythrinus* revealed similar $L50_{land}$ for the three codends
295 with overlap of their 95% CI (Table 3). Landing probability increased (>0.15) at sizes >10 cm
296 for the 40D and >12 cm for the 50D and 40S (Fig. 5). Fisher landing probability did not differ
297 between the 40D and 40S, whereas significant lower values were detected for the 50D than
298 40D or the 50D than 40S at lengths 14 - 17 cm and 14 - 16 cm, respectively (Fig. 6).

299

300 3.5. *Lophius budegassa*

301 The parameters and statistics for the best overall selection model of *L. budegassa* appear
302 in Table 3. The model generally fitted the data well, although the very low number of
303 individuals in the small sizes (<10 cm) resulted in wide 95% CI for the escape curve and the
304 left side of the bell-shaped discard curve (Fig. 5).

305 The gear selection parameter of *L. budegassa* was very low and similar among the three
306 codends (Table 3). The escape probability decreased notably (<0.3) in all cases at lengths ≥ 7
307 cm (Fig. 5) with no significant differences among the three codends (Fig. 7).

308 The discard probability of *L. budegassa* showed very high values (from 0.5 to 1.0 for sizes
309 between 5 and 22 cm) for all codends TL (Fig. 5), indicating that a large part of the total
310 amount of *L. budegassa* entering the tested gears is discarded. A significantly lower discard
311 probability was predicted for the 40D compared to 40S and 50D codends for the sizes 17-18
312 cm and 17 - 23 cm, respectively (Fig. 7). No significant differences were detected between

313 the 40S and 50D (Fig. 7). Similar indicators were obvious among the three codends with
314 overlap in their 95% CI (Table 2).

315 Landing probability of *L. budegassa* increased (>0.15) at sizes >15 cm for the 40D
316 and >18 cm for the 50D and 40S, respectively (Fig. 5). $L50_{land}$ for the 40D presented a
317 significantly lower value compared to the other codends. A significantly higher landing
318 probability for the 40D than the 40S or 50D was found at lengths from 16 to 19 cm and 17 to
319 23 cm, respectively (Fig. 7). Similar was the value of $L50_{land}$ for the 40S and 50D (Table 3).
320

321 **4. Discussion**

322 The overall size-selection during trawl fishing, based on the two sequential selection
323 processes, by the codend in the sea and by the fisher onboard the fishing vessel, was
324 modelled in the present work for four commercially important species. The model, introduced
325 by Mytilineou et al. (2018), fitted the data well in all cases in this study and simultaneously
326 predicted the escape, discard, and landing probability of the four species using only the data
327 from a selectivity experiment. In the past, these estimates were usually obtained by separate
328 studies. The results also showed that the model can be applied to more species than those
329 examined by Mytilineou et al. (2018). Nevertheless, it should be mentioned that 95% CI may
330 be wide in case of poor data.

331 The results for *M. barbatus* showed that the 40S codend was more selective than the two
332 diamond codends. $L50_{gear}$ for the 40D was significantly lower than the MCRS of 11 cm TL,
333 showing the inadequacy of this codend for the sustainability of the species. $L50_{gear}$ for the
334 40S was slightly higher than MCRS and similar to the length at first maturity of *M. barbatus*
335 ($L50_{mat}$: 12.9 cm TL; Tsikliras and Stergiou, 2013). MCRS was included between the 95% CI
336 of $L50_{gear}$ for the 50D. However, higher SR_{gear} was estimated for 50D compared to 40S,
337 resulting in i) the retention of undersized individuals (<11 cm) that will be discarded or

338 landed (in the case of 40S they escape) and ii) the escapement of individuals (≥ 17 cm TL)
339 much larger than the MCRS and the $L50_{mat}$. The fisher behaviour for this species was
340 characterised by the selection of individuals less than the MCRS as landings, depending on
341 the availability of these individuals in the catch. As a result, the landing probability was
342 higher for both the 40D and 50D than the 40S in the undersized individuals of *M. barbatus*
343 and the parameter $L50_{land}$ of the diamond codends was significantly lower than that of the
344 40S, although for the 50D it was close to the MCRS of the species. $L50_{land}$ for the 40S was
345 similar to $L50_{mat}$, indicating that the fisher selection pattern is guided towards more
346 sustainable behaviour for the stock and higher compliance with the rules when the 40S is in
347 use. The significantly lower landing probability between the 40S and 40D or 50D indicated
348 economic losses at sizes slightly above the MCRS. In contrast, higher economic losses are
349 expected using the 50D, as larger and more economically valuable individuals escape through
350 this codend. Moreover, the discard probability was very low in all cases, except for sizes
351 close to the MCRS, being higher for both diamond codends. Based on the above results, it
352 could be suggested that among the tested codends, the 40S is the most adequate gear for *M.*
353 *barbatus* in terms of the sustainability of the stock, with i) a gear selection close to $L50_{mat}$, ii)
354 lawful catch with negligible discards and iii) fisher selection pattern associated with more
355 compliance and the least economic losses in the short term. Furthermore, the 40S codend is
356 also more promising regarding the Commission Delegated Regulation (Eu) 2017/86 that
357 permits discards up to 6% of the total annual landings of *M. barbatus*.

358 The model for *M. surmuletus* showed that the 40S codend was more selective than the
359 two diamond codends, as in the case of the congeneric species *M. barbatus*. $L50_{gear}$ for the
360 40D was significantly lower than the MCRS of 11 cm TL indicating that this codend is
361 inappropriate for this species. For the 40S, $L50_{gear}$ was a little higher than MCRS, but lower
362 than the length at first maturity of *M. surmuletus* ($L50_{mat}$: 15.5 cm TL; Tsikliras and Stergiou,

363 2013). The MCRS was included between the 95% CI of the gear selection parameter for the
364 50D codend, but it was also lower than $L50_{mat}$. Moreover, the higher SR_{gear} of this codend
365 compared to the 40S had as a result the retention of more undersized individuals (<11 cm)
366 that will be discarded or landed. The fisher behaviour for this species was also characterised
367 by the selection of individuals less than the MCRS as landings, depending on the availability
368 of these individuals in the catch. Therefore, the landing probability for the undersized catch
369 of *M. surmuletus* was higher when the 40D was in use. The $L50_{land}$ of the 50D codend had a
370 similar value to MCRS, but lower than the $L50_{mat}$ of the species. Although still low, $L50_{land}$ of
371 the 40S was slightly closer to $L50_{mat}$ indicating that, fisher selection pattern would be directed
372 to a more sustainable practice for the stock in this case. Furthermore, less economic losses for
373 the fisher are expected with the 40S compared to the 50D, because the latter permit the
374 escapement of much larger than the MCRS and the $L50_{mat}$ individuals that are marketable and
375 of high economic value. Furthermore, in contrast to the 40S, the diamond codends presented
376 a higher discard probability in the sizes of juveniles. All the above let us suggest that among
377 the three codends, although none achieved $L50_{mat}$, the 40S is the most effective for *M.*
378 *surmuletus* sustainability, with a lawful catch and a fisher selection pattern associated with
379 better compliance to rules, no discards and thus less time spent by the crew in sorting, and the
380 lowest possible economic losses. Sola and Maynou (2018) tested the use of a panel with 90°
381 turned meshes in the extension part of the trawl, however, with economic losses for the
382 fishers.

383 For *P. erythrinus*, the 50D codend showed the highest $L50_{gear}$, which however cannot be
384 considered significantly different from that of the 40S, because of the overlap of their 95%
385 CI. The 40D presented the worst $L50_{gear}$ without significant difference from 40S. Thus, none
386 of the codends displayed a gear selection close to the MCRS of the species (15 cm TL),
387 which is close to the length at first maturity of the species ($L50_{mat}$: 16.4 cm TL; Tsikliras and

388 Stergiou, 2013). Furthermore, the discard likelihood did not show important differences
389 among the examined codends. Fisher selection behaviour was also similar for all tested
390 codends and was always below the MCRS and the length at first maturity. However, because
391 of a higher SR_{gear} , the 50D codend presented a higher escape probability around the MCRS
392 and the length at first maturity of the species, which might indicate a more promising pattern
393 (although not sufficiently successful) than the 40D and 40S. This means that the use of the
394 50D codend may produce some economic losses in the short term. These losses will be higher
395 in the case of a potential increase of the codend mesh size to improve gear selection and
396 reach the MCRS. Such an improvement in gear selection seems difficult unless an innovative
397 modification of the gear achieves this goal. The use of the 50D or 40S and the protection of
398 the species nursery grounds may be an alternative measure for the sustainability of the
399 species stocks and the mitigation of discards, as proposed for other species (Khoukh and
400 Maynou, 2018; Russo et al., 2019; Mytilineou et al., 2020).

401 *L. budegassa* gear selection was very negligible in all cases. Almost all individuals were
402 retained and the greatest part of the catch with sizes <22 cm has been predicted as discards.
403 Even the fish sorted as landings by the fisher were in their majority of much smaller length
404 than the length at first maturity of the species ($L50_{mat}$: 48 - 59 cm TL; Duarte et al., 2001;
405 Colmenero et al., 2013). Therefore, none of the three codends is adequate for this species in
406 terms of juvenile protection and discards. This fact is probably related to the body features of
407 this species characterized by a large head and a benthic and relatively inactive behaviour
408 inside the trawl codend (Mytilineou, unpublished data) that reduces its escape probability.
409 Gear selectivity needs considerable improvement for this species. However, considering the
410 difference between the gear selection and the $L50_{mat}$, this seems impossible without a huge
411 increase of the codend mesh size (probably resulting in the loss of other commercially
412 important catch), another innovative change in the gear design (as proposed for other species

413 in ICES WKING, 2020) or the protection of the species nursery grounds as proposed for
414 other species (Khoukh and Maynou, 2018; Russo et al., 2019; Mytilineou et al., 2020).

415 In summary, the 40D mesh in the trawl codend can be considered a mesh of low
416 selectivity, unsafe and inappropriate for the protection of juveniles, the mitigation of discards
417 and the sustainability of the stocks as also suggested by many researchers. Even in the case of
418 economic losses from the change of the 40D to another codend, the losses are mainly
419 associated with undersized, below the MCRS or the length at first maturity, catch. The 40S
420 codend was more suitable in terms of stock sustainability and with less economic losses for
421 the fisher than the 50D for the two *Mullus* species (although not reaching the MCRS for *M.*
422 *surmuletus*). No one of the 40S or 50D codends was suitable for *P. erythrinus*, although 50D
423 showed a little higher escape probability at sizes around MCRS, accompanied however by
424 more economic losses for the fisher. All the tested codends seemed harmful for *L. budegassa*.

425 The results of L_{50gear} for the studied species published in the literature are presented in
426 Table 4. Comparisons of L_{50gear} among the various researchers are not easy, because several
427 factors such as the net material, the nominal or actual mesh size, the number of mesh sizes in
428 the codend circumference, the knotted or knotless design, the catch size and shape and other
429 factors may affect the codend selectivity (e.g. Herrmann, 2005; Sala and Lucchetti, 2010).
430 Nevertheless, some of the published L_{50gear} are in agreement with our results, especially when
431 the net characteristics seemed similar (Table 4, e.g. Sala et al. 2015: for *M. barbatus*; Ordines
432 et al. 2006: for *M. surmuletus* in 40S codend; Ateş et al. 2015: for *P. erythrinus* in 40S
433 codend). No published work on the gear selection of *L. budegassa* is known. The study of
434 Tosunoğlu et al. (2008) on the 50D codend selection for the congeneric species *Lophius*
435 *piscatorius* revealed no escapees and only retained individuals, results that are similar to the
436 current study. Furthermore, based on the results of Table 4, it is worth mentioning that
437 comparing the codends with similar meshes but with different circumference for the same

438 species, in most cases, the lower the number of meshes in the codend circumference the
439 higher the L_{50gear} . Moreover, it is clear that in most of cases, L_{50gear} for the 40S codend is
440 higher than that of the 40D and 50D for *Mullus* species (Table 4). In contrast, in most cases,
441 L_{50gear} of *P. erythrinus* for the 50D was higher than that of the 40S and 40D, a fact probably
442 associated with the body shape of this species, being noticeably deep and compressed; the
443 widthwise stretching of the 50D meshes seems to benefit the escapement of this species.
444 However, the MCRS of *P. erythrinus* was achieved only when the 50D was combined with a
445 low number of meshes in the circumference (Table 4); lower than that used in commercial
446 fishery.

447 Considering the results for $L_{50fisher/discard}$, derived from the applied model and selectivity
448 data with those in the literature derived from observers onboard fishing vessels, it is worth
449 noting that these were quite comparable in the case of *M. barbatus* and *P. erythrinus* (Table
450 4). The lower values found by Damalas et al. (2018) for *M. barbatus* and *L. budegassa* and
451 Machias et al. (2004) for *M. surmuletus* may indicate a lower availability of small individuals
452 in the catch and an increase in the fisher selection behaviour nowadays, probably because of
453 the improved selectivity of the trawl codend according to the EC Regulation 1967/2006,
454 implemented years later than 2006. It should also be mentioned that the hauls conducted in
455 the present study for *L. budegassa* may not be spatially and temporally the most appropriate,
456 a fact that may have affected the population structure of the species compared to that from the
457 commercial fishery.

458 The model applied in this work was proved again to be a useful, cost-efficient approach in
459 collecting information for fisheries management, as it simultaneously predicts important
460 information on escapees, discards and fisher behaviour based on selectivity data. Moreover,
461 discards and fisher behaviour related predictions, based on one vessel and one period data,
462 were generally in accordance with those in the literature estimated from fleet-based data,

463 which supports further the applicability of the model. Concerning the codend mesh, it could
464 be suggested that the 40S codend, although not so adequate in all cases, is the most
465 sustainable compared to the 50D for the Mediterranean trawl multispecies fishery. This
466 information is useful in fisheries management since the use of the 50D is an alternative of the
467 40S according to the Council Regulation (EC) No 1967/2006. Nevertheless, within the
468 concept of the ecosystem approach to fishery management, it seems that more changes should
469 be investigated to improve the selectivity of the trawl codend with innovative gears (Brčić et
470 al., 2015) or measures (Santiago et al., 2015) along with the protection of nursery grounds,
471 particularly for species for which selectivity improvement cannot be achieved without
472 important reduction of other commercial catch and consequently fisher income. More similar
473 studies should be conducted in the future for other species since the trawl fishery in the
474 Mediterranean is a multi-species fishery.

475

476 **5. Conclusions**

477 The model applied in this work, representing the overall selectivity on a population entering
478 the trawl codend, is a cost-effective approach to collect information on the escapees, discards
479 and landings of *M. barbatus*, *M. surmuletus*, *P. erythrinus* and *L. budegassa*. The 40 mm
480 diamond mesh codend was always inappropriate for the stocks. The 40 mm square mesh
481 codend was the most effective for the sustainability of both *Mullus* species. None of the
482 codends was adequate for *P. erythrinus* and *L. budegassa*. The 50 mm diamond codend does
483 not meet the requirements of the current legislation for the Mediterranean bottom trawl in
484 terms of better selectivity compared to the 40 mm square mesh codend.

485

486 **Acknowledgements**

487 This work is part of the first author PhD thesis at the NKUA. The authors would like to
488 express their gratitude to the HCMR staff and the fishing vessel captain and crew involved in
489 the field and data processing work of EPILEXIS project (EPAL, Metro 3.5, code: 185365).
490 The authors are also grateful for the Editor's and the two anonymous Reviewers' constructive
491 comments that provided added value to this work.

492

493 **References**

494 Akaike H. 1974. A new look at the statistical model identification. *IEEE Transactions on*
495 *Automatic Control* 19, 716-722.

496 Ateş C., Deval M.C., Bök T., Tosunolu Z. 2010. Selectivity of diamond (PA) and square (PE)
497 mesh codends for commercially important fish species in the Antalya Bay, eastern
498 Mediterranean. *J. Appl. Ichthyol.* 26, 465–471.

499 Aydın C., Tokaç A., Ulaş A., Maktay B., Şensurat T. 2011. Selectivity of 40 mm square and
500 50 mm diamond mesh codends for five species in the Eastern Mediterranean demersal
501 trawl fishery. *Afr. J. Biotech.* 10 (25), 5037-5047.

502 Bellido J. M., García-Rodríguez M., García-Jiménez T., González-Aguilar M., Carbonell-
503 Quetglas A., 2017. Could the obligation to land undersized individuals increase the black
504 market for juveniles: evidence from the Mediterranean? *Fish Fish.* 18, 185–194. doi:
505 10.1111/faf.12166

506 Brčić J., Herrmann B., De Carlo F., Sala A. 2015. Selective characteristics of a shark-
507 excluding grid device in a Mediterranean trawl. *Fish. Res.* 172, 352-360.

508 Brčić J., Herrmann B., Sala A. 2018. Can a square-mesh panel inserted in front of the cod end
509 improve size and species selectivity in Mediterranean trawl fisheries? *Can. J. Fish.*
510 *Aquat. Sci.* 75, 704–713.

511 Colmenero A.I., Tuset V.M., Recasens L., Sánchez P. 2013. Reproductive biology of Black
512 Anglerfish (*Lophius budegassa*) in the northwestern Mediterranean Sea. Fish. Bull. 111,
513 390-401.

514 Commission Delegated Regulation (EU) 2017/86 of 20 October 2016 establishing a discard
515 plan for certain demersal fisheries in the Mediterranean Sea. [https://eur-](https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A32017R0086)
516 [lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A32017R0086](https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A32017R0086)

517 Council Regulation (EC) No 1626/94 of 27 June 1994 laying down certain technical
518 measures for the conservation of fishery resources in the Mediterranean.

519 Council Regulation (EC) No 1967/2006 of 21 December 2006, concerning management
520 measures for the sustainable exploitation of fishery resources in the Mediterranean Sea,
521 amending Regulation (EEC) no 2847/93 and repealing Regulation (EC) no 1626/94.,
522 2006. Off. J. Eur. Union L 409.

523 Damalas D., Ligas A., Tsagarakis K., Vassilopoulou V., Stergiou K., Kallianiotis A., Sbrana
524 M., Maynou F. 2018. The “discard problem” in Mediterranean fisheries, in the face of the
525 European Union landing obligation: the case of bottom trawl fishery and implications for
526 management. Med. Mar. Sci. 19 (3), 459-476.

527 Damalas D., Vasilopoulou V. 2013. Slack regulation compliance in the Mediterranean
528 fisheries: a paradigm from the Greek Aegean Sea demersal trawl fishery, modelling
529 discard ogives. Fish. Manag. Ecol. 20, 21-33.

530 Dereli H., Aydin C. 2016. Selectivity of Commercial and Alternative Codends for Four
531 Species in the Eastern Mediterranean Demersal Trawl Fishery. Turk. J. Fish. Aquat. Sc.
532 16, 971-992.

533 Duarte R., Azevedoa M., Landab J., Pereda P. 2001. Reproduction of anglerfish (*Lophius*
534 *budegassa* Spinola and *Lophius piscatorius* Linnaeus) from the Atlantic Iberian coast.
535 Fish. Res. 53, 349 - 361

536 Efron B. 1982. The jackknife, the bootstrap and other resampling plans. SIAM Monograph
537 No. 38, CBSM-NSF.

538 FAO 2020. The State of Mediterranean and Black Sea Fisheries 2020. General Fisheries
539 Commission for the Mediterranean. Rome. <https://doi.org/10.4060/172> pp.

540 Frandsen R.P., Madsen N., Krag L.A. 2010. Selectivity and escapement of five commercial
541 fishery species in standard square- and diamond-mesh codends. ICES J. Mar. Sci. 67,
542 1721-1731.

543 Fryer R. 1991. A model of the between-haul variation in selectivity. ICES J. Mar. Sci. 48,
544 281–290.

545 Herrmann B. 2005. Effect of catch size and shape on the selectivity of diamond mesh cod-
546 ends I. Model development. Fish. Res. 71, 1-13.

547 Herrmann B., Mieske B., Stepputtis D., Krag L.A., Madsen N., Noack T. 2013. Modelling
548 towing and haul-back escape patterns during the fishing process: a case study for cod,
549 plaice, and flounder in the demersal Baltic Sea cod fishery. ICES J. Mar. Sci. 70 (4), 850-
550 863. doi: 10.1093/icesjms/fst032.

551 Herrmann B., Sistiaga M., Larsen R.B., Brinkhof J. 2019. Effect of three different codend
552 designs on the size selectivity of juvenile cod in the Barents Sea shrimp trawl fishery.
553 Fish. Res. DOI: [10.1016/j.fishres.2019.105337](https://doi.org/10.1016/j.fishres.2019.105337)

554 Herrmann B., Sistiaga M., Nielsen K.N., Larsen R.B. 2012. Understanding the size selectivity
555 of redfish (*Sebastes* spp.) in North Atlantic trawl codends. J. Northw. Atl. Fish. Sci. 44,
556 1–13.

557 ICES WKING 2020. Workshop on Innovative Fishing Gears (WKING), May-September
558 2020, 113pp. (in press)

559 Khoukh M., Maynou F. 2018. Spatial management of the European hake *Merluccius*
560 *merluccius* fishery in the Catalan Mediterranean: Simulation of management alternatives
561 with the InVEST model. *Sci. Mar.* 82S1: 175-188.
562 <https://doi.org/10.3989/scimar.04748.18A>

563 Larsen R.B., Sistiaga M., Herrmann B., Brinkhof J., Tatone I., Santos J. 2018. The effect of
564 Nordmøre grid length and 1 angle on codend entry of bycatch fish species and shrimp
565 catches. *Can. J. Fish. Aquat. Sci.* 76 (2), 308-319.
566 <https://doi.org/10.1139/cjfas-2018-0069>

567 Machias A., Vassilopoulou V., Vatsos D., Bekas P., Kallianotis A., Papaconstantinou C. et al.
568 2004. Bottom trawl discards in the northeastern Mediterranean Sea. *Fish. Res.* 53, 181–
569 195. doi: 10.1016/S0165-7836(00) 00298-8

570 Millar R.B. 1993. Incorporation of between-haul variation using bootstrapping and
571 nonparametric estimation of selection curves. *Fish. Bull.* 91, 564–572.

572 Mytilineou Ch., Herrmann B., Mantopoulou-Palouka D., Sala A., Megalofonou P. 2018.
573 Modelling gear and fishers size selection for escapees, discards, and landings: a case
574 study in Mediterranean trawl fisheries. *ICES J. Mar. Sci.* 75 (5), 1693-1709.

575 Mytilineou Ch., Herrmann B., Kavadas S., Smith C., Megalofonou P. 2020. Combining
576 selection models and population structures to inform fisheries management: a case study
577 on hake in the Mediterranean bottom trawl fishery. *Mediterr. Mar. Sci.* 21/2, 360-371.
578 doi:<https://doi.org/10.12681/mms.22191>

579 Ordines, F., Massuti, E., Guijarro, B., and Mas, R. 2006. Diamond vs. square mesh codend in
580 a multi-species trawl fishery of the western Mediterranean: effects on catch composition,
581 yield, size selectivity and discards. *Aquat. Liv. Res.*, 19, 329-338.

582 Özbilgin H., Eryasar A.R., Gokce G., Özbilgin Y.D., Bozaoglu A.S., Kalecik E., Herrmann B.
583 2015. Size selectivity of hand and machine woven codends and short term commercial
584 loss in the Northeastern Mediterranean. *Fish. Res.* 164, 73-85.

585 Özbilgin H., Metin G., Tosunoğlu Z., Tokaç A., Kaykaç, H., Aydın C. 2012. Seasonal
586 variation in the trawl codend selectivity of common pandora (*Pagellus erythrinus*). *J.*
587 *Appl. Ichthyol.* 28, 194-199.

588 Özbilgin H., Tosunoğlu Z. 2003. Comparison of the selectivities of double and single
589 codends. *Fish. Res.* 63, 143–147.

590 Ragheb E., Akel K., Rizkalla S.I. 2019. Analyses of the non-target catch from the Egyptian
591 Mediterranean El Sayed Haroun trawlers, off Port Said. *Egypt. J. Aqu. Res.* 45, 239-246.

592 Regulation (EU) 2019/1241 of the European Parliament and of the Council of 20 June 2019
593 on the conservation of fisheries resources and the protection of marine ecosystems
594 through technical measures.

595 Russo T., D'Andrea L., Franceschini S., Accadia P., Cucco A., Garofalo G., Gristina M.,
596 Parisi A., Quattrocchi G., Sabatella R.F., Sinerchia M., Canu D.M., Cataudella S.,
597 Fiorentino F. 2019. Simulating the Effects of Alternative Management Measures of Trawl
598 Fisheries in the Central Mediterranean Sea: Application of a Multi-Species Bio-economic
599 Modeling Approach. *Front. Mar. Sci.* 6, 542. doi: 10.3389/fmars.2019.00542

600 Sala, A., Lucchetti, A. 2010. The effect of mesh configuration and codend circumference on
601 selectivity in the Mediterranean trawl *Nephrops* fishery. *Fish. Res.* 103, 63-72.

602 Sala A., Lucchetti A., Perdichizzi A., Herrmann B., Rinelli P. 2015. Is square -mesh better
603 selective than larger mesh? A perspective on the management for Mediterranean trawl
604 fisheries. *Fish. Res.* 161, 182-190.

605 Santiago J.L., Ballesteros M.A., Chapela R., Silva C., Nielsen K.N., Rangel M., Erzini K.,
606 Wise L., Campos A., Borges M.F., Sala A., Virgili M., Viðarsson J.R., Baudron A.,

607 Fernandes P.G. 2015. Is Europe ready for a results-based approach to fisheries
608 management? The voice of stakeholders. *Mar. Pol.* 56, 86-97.

609 Sistiaga M., Herrmann B., Grimaldo E., Larsen R., 2010. Assessment of dual selection in grid
610 based selectivity systems. *Fish. Res.* 105, 187–199.

611 Sola I., Maynou F. 2018. Assessment of the relative catch performance of hake, red mullet
612 and striped red mullet in a modified trawl extension with T90 netting. *Sci. Mar.* 82S1:
613 000-000. <https://doi.org/10.3989/scimar.04711.04A>

614 Tokaç A., Herrmann B., Aydın C., Kaykac H., Ünlüler A., Gökc G. 2014. Predictive models
615 and comparison of the selectivity of standard (T0) and turned mesh (T90) codends for
616 three species in the Eastern Mediterranean. *Fish. Res.* 150, 76– 88.

617 Tosunoğlu Z., Aydın C., Özyayın O. 2008. Selectivity of a 50-mm diamond mesh knotless
618 polyethylene codend for commercially important fish species in the Aegean Sea. *J. Appl.*
619 *Ichthyol.* 24, 311–315.

620 Tosunoğlu Z., Doğanıyılmaz Özbilgin Y., Özbilgin, H. 2003. Body shape and trawl cod end
621 selectivity for nine commercial fish species. *J. Mar. Biol. Ass. U.K.* 83, 1309-1313.

622 Tsagarakis K., Carbonell A., Brčić J., Bellido J.M., Carbonara P., Casciaro L., Edridge A. et
623 al. 2017. Old info for a New Fisheries Policy: Discard Ratios and Lengths at Discarding
624 in EU Mediterranean Bottom Trawl Fisheries. *Front. Mar. Sci.* 4, 99.
625 <https://doi.org/10.3389/fmars.2017.00099>.

626 Tsikliras A. and Stergiou K. 2013. Size at maturity of Mediterranean marine fishes. *Rev Fish*
627 *Biol Fisher.* 24, 219–268. doi: 10.1007/s11160-013-9330-x

628 Vasilakopoulos P., O'Neill F.G., Marsall T. 2015. The unfulfilled potential of fisheries
629 selectivity to promote sustainability. *Fish and Fisheries*, doi:10.1111/faf.12117.

630 Wileman D., Ferro R.S.T., Fonteyne R., Millar R.B. 1996. Manual of methods of measuring
631 the selectivity of towed fishing gear. ICES Coop. Res. Report No. 215, 126 pp.

632 **Figures captions**

633

634 **Fig. 1.** Map of the study area, where the hauls (red diamonds) of the experimental fishing
635 were conducted; isobath 50 m: dots line, isobath 100 m: continuous line, isobath 300 m:
636 dashed line.

637

638 **Fig. 2.** Size-selection curves for *Mullus barbatus* and *Mullus surmuletus* when using 40 mm
639 diamond mesh (40D), 40 mm square mesh (40S) or 50 mm diamond mesh (50D) in the trawl
640 codend. Blue curves and triangles: gear escape probability (p_{esc}) and associated experimental
641 ratios; grey curves and crosses: discard probability (p_{disc}) and associated experimental ratios;
642 red curves and dots: landing probability (p_{land}) and associated experimental ratios. Coloured
643 areas around the curves: Efron percentile 95% confidence intervals.

644

645 **Fig. 3.** Difference in the *Mullus barbatus* size-dependent escape (E), discard (D) and landing
646 probability (L) (in blue, grey and red colour, respectively) between 40D - 40S, 50D - 40D and
647 50D - 40S codends. 40D, 40S, 50D: trawl codend with 40 mm diamond mesh, 40 mm square
648 mesh, 50 mm diamond mesh, respectively; coloured areas around lines: 95% Efron percentile
649 confidence intervals.

650

651 **Fig. 4.** Difference in the *Mullus surmuletus* size-dependent escape (E), discard (D) and
652 landing probability (L) (in blue, grey and red colour, respectively) between 40D - 40S, 50D -
653 40D and 50D - 40S codends. 40D, 40S, 50D: trawl codend with 40 mm diamond mesh, 40
654 mm square mesh, 50 mm diamond mesh, respectively; coloured areas around lines: 95%
655 Efron percentile confidence intervals.

656


657 **Fig. 5.** Size-selection curves for *Pagellus erythrinus* and *Lophius budegassa* when using 40
658 mm diamond mesh (40D), 40 mm square mesh (40S) and 50 mm diamond mesh (50D) in the
659 trawl codend. Blue curves and triangles: gear escape probability (p_{esc}) and associated
660 experimental ratios; grey curves and crosses: discard probability (p_{disc}) and associated
661 experimental ratios; red curves and dots: landing probability (p_{land}) and associated
662 experimental ratios. Coloured areas around the curves: Efron percentile 95% confidence
663 intervals.


664


665 **Fig. 6.** Difference in the *Pagellus erythrinus* size-dependent escape (E), discard (D) and
666 landing probability (L) (in blue, grey and red colour, respectively) between 40D - 40S, 50D -
667 40D and 50D – 40S codends. 40D, 40S, 50D: trawl codend with 40 mm diamond mesh, 40
668 mm square mesh, 50 mm diamond mesh, respectively; coloured areas around lines: 95%
669 Efron percentile confidence intervals.


670


671 **Fig. 7.** Difference in the *Lophius budegassa* size-dependent escape (E), discard (D) and
672 landing probability (L) (in blue, grey and red colour, respectively) between 40D - 40S, 50D -
673 40D and 50D – 40S codends. 40D, 40S, 50D: trawl codend with 40 mm diamond mesh, 40
674 mm square mesh, 50 mm diamond mesh, respectively; coloured areas around lines: 95%
675 Efron percentile confidence intervals.




















Table 1

Selectivity parameters for *Mullus barbatus* and *Mullus surmuletus* for the overall selection model describing the size-selectivity of the gear ($L50_{gear}$, SR_{gear} , $1/\delta_{gear}$), the fisher size-selection ($L50_{fisher}$, SR_{fisher} , $1/\delta_{fisher}$) and the landing probability ($L50_{land}$, SR_{land}) in the trawl codend when using the 40D (40 mm diamond), 40S (40 mm square) or 50D (50 mm diamond) mesh; 95% confidence intervals (*Efron percentile*) are shown in parenthesis; $1/\delta$ is presented in the case of Richard model. (G: gear selectivity model; F: fisher selection model; DOF: degrees of freedom; AIC: Akaike criterion).

Species	Model Parameter	Codend		
		40D	40S	50D
<i>Mullus barbatus</i>		Model		
		G: Richard F: Gompertz	G: Richard F: Logit	G: Gompertz F: Probit
	$L50_{gear}$	9.34 (7.77-10.04)	13.31 (12.36-13.68)	10.83 (7.94-12.13)
	SR_{gear}	2.00 (0.47-2.49)	2.23 (1.83-3.97)	4.73 (3.44-7.19)
	$1/\delta_{gear}$	0.58 (0.10-10.00)	0.56 (0.15-1.40)	
	$L50_{fisher}$	8.71 (0.10-9.45)	10.48 (0.10-11.73)	9.41 (0.10-10.34)
	SR_{fisher}	1.53 (1.04-4.08)	0.10 (0.10-0.10)	1.67 (0.10-2.29)
	$L50_{land}$	9.80 (7.94-10.26)	13.31 (12.36-13.68)	11.19 (10.15-12.18)
	SR_{land}	1.61 (0.59-1.91)	2.23 (1.83-3.97)	3.60 (2.27-5.14)
	p-value	0.2169	0.2589	0.3711
	Deviance	43.41	44.27	38.17
	DOF	37	39	36
AIC	421.87	2668.21	2248.49	
<i>Mullus surmuletus</i>		Model		
		G: Probit F: Probit	G: Logit F: Logit	G: Gompertz F: Gompertz
	$L50_{gear}$	8.40 (0.10-9.69)	12.04 (11.37-12.63)	10.84 (9.02-12.35)
	SR_{gear}	2.20 (0.10-4.72)	1.65 (1.17-2.05)	5.81 (2.49-8.42)
	$L50_{fisher}$	8.05 (4.10-9.90)	-	9.08 (0.10-10.27)
	SR_{fisher}	2.10 (0.20-3.84)	-	1.09 (0.10-1.82)
	$L50_{land}$	9.10 (5.69-10.16)	12.04 (11.37-12.63)	11.06 (9.55-12.45)
	SR_{land}	1.78 (0.47-3.54)	1.65 (1.17-2.05)	4.61 (2.33- 5.94)
	p-value	1.0000	1.0000	0.9998
	Deviance	7.47	2.72	9.98
	DOF	40	38	30
	AIC	136.19	211.46	486.08

Table 2

Discard parameters (with confidence intervals) based on the best model for the overall size-selection process by the gear and the fisher; ($DR_{0.05}, DR_{0.25}, DR_{0.5}, DR_{0.75}, DR_{0.95}$): discard range (cm) at several probability levels, Dp_{max} : maximum discard probability value, LDp_{max} : length (cm) at the maximum discard probability and $DA_{0.05}$: surface of the discard bell-shaped curve when probability ≥ 0.05 .

Species	Parameter	CODEND		
		40D	40S	50D
<i>Mullus barbatus</i>	$DR_{0.05}$	5.05 (0.00-6.61)	0.49 (0.00-7.02)	4.68 (0.00-9.81)
	$DR_{0.25}$	0.00 (0.00-2.12)	0.00 (0.00-0.20)	0.00 (0.00-5.05)
	$DR_{0.5}$	0.00 (0.00-0.00)	0.00 (0.00-0.00)	0.00 (0.00-0.00)
	$DR_{0.75}$	0.00 (0.00-0.00)	0.00 (0.00-0.00)	0.00 (0.00-0.00)
	$DR_{0.95}$	0.00 (0.00-0.00)	0.00 (0.00-0.00)	0.00 (0.00-0.00)
	Dp_{max}	0.17 (0.01-0.39)	0.06 (0.00-0.27)	0.18 (0.00-0.46)
	LDp_{max}	8.65 (7.57-10.77)	9.83 (0.00-10.73)	8.87 (0.00-9.64)
	$DA_{0.05}$	0.57 (0.00-1.16)	0.00 (0.00-0.93)	0.55 (0.00-2.56)
<i>Mullus surmuletus</i>	$DR_{0.05}$	4.93 (1.32-10.60)	0.00 (0.00-0.00)	5.04 (0.00-8.80)
	$DR_{0.25}$	0.00 (0.00-7.79)	0.00 (0.00-0.00)	0.00 (0.00-3.22)
	$DR_{0.5}$	0.00 (0.00-5.83)	0.00 (0.00-0.00)	0.00 (0.00-0.00)
	$DR_{0.75}$	0.00 (0.00-3.41)	0.00 (0.00-0.00)	0.00 (0.00-0.00)
	$DR_{0.95}$	0.00 (0.00-0.35)	0.00 (0.00-0.00)	0.00 (0.00-0.00)
	Dp_{max}	0.21 (0.06-1.00)	0.00 (0.00-0.00)	0.22 (0.00-0.40)
	LDp_{max}	8.15 (1.47-9.36)	0.00 (0.00-0.00)	8.14 (0.00-9.87)
	$DA_{0.05}$	0.66 (0.07-3.41)	0.00 (0.00-0.00)	0.66 (0.00-1.79)
<i>Pagellus erythrinus</i>	$DR_{0.05}$	10.59 (6.00-15.29)	6.18 (0.00-7.00)	7.11 (1.95-13.98)
	$DR_{0.25}$	4.95 (3.60-13.08)	3.49 (0.00-4.49)	0.23 (0.00-5.21)
	$DR_{0.5}$	2.16 (1.19-11.46)	0.69 (0.00-2.64)	0.00 (0.00-0.35)
	$DR_{0.75}$	0.00 (0.00-9.78)	0.00 (0.00-0.00)	0.00 (0.00-0.00)
	$DR_{0.95}$	0.00 (0.00-7.37)	0.00 (0.00-0.00)	0.00 (0.00-0.00)
	Dp_{max}	0.65 (0.56-1.00)	0.54 (0.02-0.72)	0.26 (0.07-0.49)
	LDp_{max}	11.00 (1.50-11.97)	11.78 (11.29-13.43)	11.76 (10.99-12.73)
	$DA_{0.05}$	2.99 (2.16-8.37)	1.80 (0.00-2.66)	1.00 (0.12-3.22)
<i>Lophius budegassa</i>	$DR_{0.05}$	21.02 (17.13-24.61)	24.96 (13.59-26.26)	25.96 (20.22-31.85)
	$DR_{0.25}$	15.61 (12.81-20.92)	21.93 (11.43-24.25)	20.57 (16.28-27.24)
	$DR_{0.5}$	12.36 (10.12-19.13)	17.82 (9.71-23.05)	16.83 (13.63-23.93)
	$DR_{0.75}$	9.42 (7.59-17.38)	13.73 (7.64-22.52)	13.12 (10.66-20.69)
	$DR_{0.95}$	5.51 (4.01-15.69)	7.78 (2.27-22.10)	7.81 (5.75-17.08)
	Dp_{max}	1.00 (1.00-1.00)	1.00 (0.96-1.00)	1.00 (0.99-1.00)
	LDp_{max}	12.65 (1.50-13.51)	16.83 (1.50-22.50)	12.44 (1.50-14.14)
	$DA_{0.05}$	12.55 (10.24-18.86)	17.42 (9.67-21.33)	16.71 (13.55-22.81)

Table 3

Selectivity parameters for *Pagellus erythrinus* and *Lophius budegassa* for the overall selection model describing the size-selectivity of the gear ($L50_{gear}, SR_{gear}, 1/\delta_{gear}$), the fisher size-selection ($L50_{fisher}, SR_{fisher}, 1/\delta_{fisher}$) and the landing probability ($L50_{land}, SR_{land}$) in the trawl codend when using the 40D (40 mm diamond), 40S (40 mm square) or 50D (50 mm diamond) mesh; 95% confidence intervals (*Efron percentile*) are shown in parenthesis; $1/\delta$ is presented in the case of Richard model. (G: gear selectivity model; F: fisher selection model; DOF: degrees of freedom; AIC: Akaike criterion).

Species	Model	Codend		
		40D	40S	50D
<i>Pagellus erythrinus</i>	Parameter	G: Richard F: Logit	G: Gompertz F: Logit	G: Richard F: Gompertz
	$L50_{gear}$	9.72 (0.10-10.75)	11.02 (10.38-12.79)	13.40 (11.71-13.94)
	SR_{gear}	2.69 (0.10-7.66)	2.39 (1.40-2.73)	3.43 (2.00-6.92)
	$1/\delta_{gear}$	0.10 (0.10-10.00)		0.29 (0.10-10.00)
	$L50_{fisher}$	12.07 (11.48-13.31)	12.81 (0.10-13.34)	12.62 (11.66-13.13)
	SR_{fisher}	1.86 (1.31-2.68)	1.47 (0.88-6.15)	1.04 (0.10-1.55)
	$L50_{land}$	12.08 (11.49-13.31)	13.08 (12.41-13.45)	13.78 (12.80-14.21)
	SR_{land}	1.74 (1.31-2.68)	1.54 (0.97-2.06)	2.09 (1.59-2.63)
	p-value	1.0000	0.2350	0.9957
	Deviance	17.04	46.09	18.32
	DOF	47	40	37
	AIC	695.58	1048.59	499.78
	<i>Lophius budegassa</i>	Parameter	G: Probit F: Gompertz	G: Probit F: Gompertz
$L50_{gear}$		4.71 (0.10-6.57)	4.43 (0.10-11.97)	5.27 (0.10-7.98)
SR_{gear}		3.53 (0.10-4.69)	5.88 (0.10-9.59)	3.06 (0.10-3.40)
$L50_{fisher}$		17.06 (15.97-19.49)	22.25 (19.75-24.00)	22.10 (19.93-24.55)
SR_{fisher}		2.56 (1.22-3.49)	2.20 (0.10-3.94)	4.31 (2.18-6.51)
$L50_{land}$		17.06 (15.97-19.49)	22.25 (19.76-24.00)	22.10 (19.93-24.55)
SR_{land}		2.56 (1.22-3.49)	2.20 (0.10-3.93)	4.30 (2.18-6.51)
p-value		1.0000	1.0000	1.0000
Deviance		6.69	3.27	19.68
DOF		68	60	66
AIC		95.48	28.89	114.63

Table 4. L_{50gear} (length at which 50% of the individuals are retained by the trawl codend) and $L_{50fisher/discard}$ (length at which 50% of the retained in the codend individuals are discarded by the fisher) for *M. barbatus*, *M. surmuletus*, *P. erythrinus* and *L. budegassa* from the Mediterranean Sea published in the literature. The results of the present work are also shown. Mesh characteristics are also given.

Species	Mesh	L_{50gear} (cm TL)	$L_{50fisher/discard}$ (cm TL)	Reference	Area
<i>M. barbatus</i>	40D600_PE*	10.60		Tosunoğlu et al. (2003)	E. Aegean Sea
	40D600_PE	10.1		Özbilgin and Tosunoğlu (2003)	E. Aegean Sea
	40D220_PE	9.14			
	40T90220_PE	12.41			
	44D200_PE	11.35		Tokaç et al. (2014)	E. Aegean Sea
	44T200_PE	14.62			
	50D176_PE	14.66			
	44D200_PA	10.7		Ateş et al. (2010)	Antalya Bay (Levantine Sea)
	40S100_PE	14.2			
	50D200_PE	15.2		Aydin et al. (2011)	E. Aegean Sea
	40S100_PE	14.4			
	44D320_PA	8.58			
	44S160_PA	13.20			
	54D256_PA	11.63		Sala et al. (2015)	Tyrrhenian Sea
	54S128_PA	17.28			
	44D400_PE_handmade	7.1			
	44D300_PE_machine	8.4			
	50D265_PE_machine	12.1		Özbilgin et al. (2015)	Mersin Bay (Levantine Sea)
	40S150_PE_machine	14.1			
	44D300_PE	11.1			
	50D264_PE	12.9			
	40T330_PE	13.6		Dereli and Aydin (2016)	E. Aegean Sea
	40S165_PE	12.9			
	50D246_PE	9.81		Brčić et al. (2018)	Tyrrhenian Sea
	40D400_PA	9.34	8.71		
	50D340_PA	10.83	9.41	Present work	S. Aegean Sea
40S200_PA	13.31	10.48			
28D			10.1-11.1	Machias et al. (2004)	E. Ionian Sea
40D/40S			6.2-7.4	Damalas et al. (2018)	Aegean Sea
<i>M. surmuletus</i>	40D_PE	4.5			
	40S_PE	12.2		Ordines et al. (2006)	Balearic Isl.
	40D400_PA	8.40	8.05		
	50D340_PA	10.84	9.08	Present work	S. Aegean Sea
	40S200_PA	12.04	-		
<i>P. erythrinus</i>	40D600_PE	10.5		Özbilgin and Tosunoğlu (2003)	E. Aegean Sea
	40D600_PE	10.80		Tosunoğlu et al. (2003)	E. Aegean Sea
	40D_PE	-			
	40S_PE	10.4		Ordines et al. (2006)	Balearic Isl.
	44D200_PA	11.8			
	40S100_PE	11.0		Ateş et al. (2010)	Antalya Bay (Levantine Sea)
	50D200_PE	15.0			
	40S100_PE	13.1		Aydin et al. (2011)	E. Aegean Sea
	44D400_PE_handmade	8.3			
	44D300_PE_machine	11.7			
	50D265_PE_machine	15.1		Özbilgin et al. (2015)	Mersin Bay (Levantine Sea)
	40S150_PE_machine	13.0			
	40D220_PE	8.73			
	40T90220_PE	10.23			
	44D200_PE	11.10		Tokaç et al. (2014)	E. Aegean Sea
44T200_PE	12.76				
50D176_PE	14.66				

	40D400_PA	9.72	12.07		
	50D340_PA	13.40	12.62	Present work	S. Aegean Sea
	40S200_PA	11.02	12.81		
	28D		12.2-13.2	Machias et al. (2004)	E. Ionian Sea
	40D		~12.0	Damalas & Vassilopoulou (2013)	Aegean Sea
<i>L. budegassa</i>	40D400_PA	4.71	17.06		
	50D340_PA	5.27	22.10	Present work	S. Aegean Sea
	40S200_PA	4.43	22.25		
	40D		14.9	Damalas et al. (2018)	Aegean Sea
	40D/40S		10.2		

* Mesh description: i) mesh size in mm, ii) mesh configuration (D: diamond, S: square, T: 90° turned mesh), iii) number of meshes in codend circumference, iv) twine material (PA: polyamid, PE: polyethylen).